

Anti-USP38 Antibody
Rabbit polyclonal antibody to USP38
Catalog # AP60649

Specification

Anti-USP38 Antibody - Product Information

Application	WB
Primary Accession	O8NB14
Other Accession	O8BW70
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	116546

Anti-USP38 Antibody - Additional Information

Gene ID 84640

Other Names

KIAA1891; Ubiquitin carboxyl-terminal hydrolase 38; Deubiquitinating enzyme 38; HP43.8KD; Ubiquitin thioesterase 38; Ubiquitin-specific-processing protease 38

Target/Specificity

Recognizes endogenous levels of USP38 protein.

Dilution

WB~~WB (1/500 - 1/1000)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-USP38 Antibody - Protein Information

Name USP38

Synonyms KIAA1891

Function

Deubiquitinating enzyme that plays a role in various cellular processes, including DNA repair, cell cycle regulation, and immune response (PubMed: [22689415](http://www.uniprot.org/citations/22689415)), PubMed: [30497519](http://www.uniprot.org/citations/30497519), PubMed: [31874856](http://www.uniprot.org/citations/31874856), PubMed: [35238669](http://www.uniprot.org/citations/35238669)). Plays a role

in the inhibition of type I interferon signaling by mediating the 'Lys-33' to 'Lys-48' ubiquitination transition of TBK1 leading to its degradation (PubMed:27692986). Cleaves the ubiquitin chain from the histone demethylase LSD1/KDM1A and prevents it from degradation by the 26S proteasome, thus maintaining LSD1 protein level in cells (PubMed:30497519). Plays a role in the DNA damage response by regulating the deacetylase activity of HDAC1 (PubMed:31874856). Mechanistically, removes the 'Lys-63'-linked ubiquitin chain promoting the deacetylase activity of HDAC1 in response to DNA damage (PubMed:31874856). Acts also as a specific deubiquitinase of histone deacetylase 3/HDAC3 and cleaves its 'Lys-63'-linked ubiquitin chains to lower its histone deacetylase activity (PubMed:32404892). Regulates MYC levels and cell proliferation via antagonizing ubiquitin E3 ligase FBXW7 thereby preventing MYC 'Lys-48'- linked ubiquitination and degradation (PubMed:34102342). Participates in antiviral response by removing both 'Lys-48'-linked and 'Lys-63'- linked polyubiquitination of Zika virus envelope protein E (PubMed:34696459). Constitutively associated with IL-33R/IL1RL1, deconjugates its 'Lys-27'-linked polyubiquitination resulting in its autophagic degradation (PubMed:35238669).

Cellular Location

Cytoplasm. Nucleus Note=In response to DNA damage, recruited to DNA damage sites in the nucleus.

Tissue Location

Highly expressed in skeletal muscle. Expressed in adrenal gland.


Anti-USP38 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-USP38 Antibody - Images





Western blot analysis of USP38 expression in A549 (A), PC12 (B) whole cell lysates.

Anti-USP38 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human USP38. The exact sequence is proprietary.