

**Anti-iNOS Antibody**  
Rabbit polyclonal antibody to iNOS  
Catalog # AP60696

### Specification

---

#### Anti-iNOS Antibody - Product Information

Application	WB
Primary Accession	<a href="#">P35228</a>
Other Accession	<a href="#">P29477</a>
Reactivity	Human, Mouse, Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	131117

#### Anti-iNOS Antibody - Additional Information

Gene ID 4843

#### Other Names

NOS2A; Nitric oxide synthase, inducible; Hepatocyte NOS; HEP-NOS; Inducible NO synthase; Inducible NOS; iNOS; NOS type II; Peptidyl-cysteine S-nitrosylase NOS2

#### Target/Specificity

Recognizes endogenous levels of iNOS protein.

#### Dilution

WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200)

#### Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

#### Storage

Store at -20 °C. Stable for 12 months from date of receipt

#### Anti-iNOS Antibody - Protein Information

Name NOS2 ([HGNC:7873](#))

Synonyms NOS2A

#### Function

Produces nitric oxide (NO) which is a messenger molecule with diverse functions throughout the body (PubMed: [7504305](http://www.uniprot.org/citations/7504305) target="\_blank">7504305</a>, PubMed: [7531687](http://www.uniprot.org/citations/7531687) target="\_blank">7531687</a>, PubMed: [7544004](http://www.uniprot.org/citations/7544004) target="\_blank">7544004</a>, PubMed: [7682706](http://www.uniprot.org/citations/7682706) target="\_blank">7682706</a>). In macrophages, NO mediates tumoricidal and bactericidal

actions. Also has nitrosylase activity and mediates cysteine S-nitrosylation of cytoplasmic target proteins such PTGS2/COX2 (By similarity). As component of the iNOS-S100A8/9 transnitrosylase complex involved in the selective inflammatory stimulus-dependent S-nitrosylation of GAPDH on 'Cys-247' implicated in regulation of the GAIT complex activity and probably multiple targets including ANXA5, EZR, MSN and VIM (PubMed:<a href="http://www.uniprot.org/citations/25417112" target="\_blank">25417112</a>). Involved in inflammation, enhances the synthesis of pro-inflammatory mediators such as IL6 and IL8 (PubMed:<a href="http://www.uniprot.org/citations/19688109" target="\_blank">19688109</a>).

#### Cellular Location

Cytoplasm, cytosol. Note=Localizes as discrete foci scattered throughout the cytosol and in the presence of SPSB1 and SPSB4, exhibits a more diffuse cytosolic localization.

#### Tissue Location


Expressed in the liver, retina, bone cells and airway epithelial cells of the lung. Not expressed in the platelets Expressed in chondrocytes (PubMed:7504305)

### Anti-iNOS Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-iNOS Antibody - Images



Western blot analysis of iNOS expression in Hela (A), A549 (B), MCF7 (C), mouse brain (D) whole cell lysates.



Immunohistochemical analysis of iNOS staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

#### **Anti-iNOS Antibody - Background**

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human iNOS. The exact sequence is proprietary.