

Anti-Recoverin Antibody
Rabbit polyclonal antibody to Recoverin
Catalog # AP60720**Specification**

Anti-Recoverin Antibody - Product Information

Application	WB, IF
Primary Accession	P35243
Other Accession	P34057
Reactivity	Human, Mouse, Rat, Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	23130

Anti-Recoverin Antibody - Additional Information**Gene ID** 5957**Other Names**

RCV1; Recoverin; Cancer-associated retinopathy protein; Protein CAR

Target/Specificity

Recognizes endogenous levels of Recoverin protein.

Dilution

WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200), IF/IC (1/100 - 1/500)

IF~~WB (1/500 - 1/1000), IH (1/100 - 1/200), IF/IC (1/100 - 1/500)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-Recoverin Antibody - Protein Information**Name** RCVRN**Synonyms** RCV1**Function**

Acts as a calcium sensor and regulates phototransduction of cone and rod photoreceptor cells (By similarity). Modulates light sensitivity of cone photoreceptor in dark and dim conditions (By similarity). In response to high Ca(2+) levels induced by low light levels, prolongs RHO/rhodopsin activation in rod photoreceptor cells by binding to and inhibiting GRK1-mediated phosphorylation of RHO/rhodopsin (By similarity). Plays a role in scotopic vision/enhances vision in dim light by enhancing signal transfer between rod photoreceptors and rod bipolar cells (By similarity).

Improves rod photoreceptor sensitivity in dim light and mediates response of rod photoreceptors to facilitate detection of change and motion in bright light (By similarity).

Cellular Location

Photoreceptor inner segment {ECO:0000250|UniProtKB:P34057}. Cell projection, cilium, photoreceptor outer segment {ECO:0000250|UniProtKB:P34057}. Photoreceptor outer segment membrane {ECO:0000250|UniProtKB:P21457}; Lipid-anchor {ECO:0000250|UniProtKB:P21457}; Cytoplasmic side {ECO:0000250|UniProtKB:P21457}. Perikaryon {ECO:0000250|UniProtKB:P34057}. Note=Primarily expressed in the inner segments of light-adapted rod photoreceptors, approximately 10% of which translocates from photoreceptor outer segments upon light stimulation (By similarity). Targeting of myristoylated protein to rod photoreceptor outer segments is calcium dependent (By similarity) {ECO:0000250|UniProtKB:P21457, ECO:0000250|UniProtKB:P34057}

Tissue Location


Retina and pineal gland.

Anti-Recoverin Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-Recoverin Antibody - Images



Western blot analysis of Recoverin expression in mouse testis (A), rat testis (B) whole cell lysates.



Immunohistochemical analysis of Recoverin staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of Recoverin staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Anti-Recoverin Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Recoverin. The exact sequence is proprietary.