

Anti-B4GALT1 Antibody
Rabbit polyclonal antibody to B4GALT1
Catalog # AP60753

Specification

Anti-B4GALT1 Antibody - Product Information

| | |
|-------------------|-----------------------------------|
| Application | WB |
| Primary Accession | P15291 |
| Other Accession | P15535 |
| Reactivity | Human, Mouse, Rat, Monkey, Bovine |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 43920 |

Anti-B4GALT1 Antibody - Additional Information

Gene ID 2683

Other Names

GGTB2; Beta-1, 4-galactosyltransferase 1; Beta-1, 4-GalTase 1; Beta4Gal-T1; b4Gal-T1; UDP-Gal:beta-GlcNAc beta-1, 4-galactosyltransferase 1; UDP-galactose:beta-N-acetylglucosamine beta-1, 4-galactosyltransferase 1

Target/Specificity

Recognizes endogenous levels of B4GALT1 protein.

Dilution

WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-B4GALT1 Antibody - Protein Information

Name B4GALT1 ([HGNC:924](#))

Synonyms GGTB2

Function

[Beta-1,4-galactosyltransferase 1]: The Golgi complex form catalyzes the production of lactose in the lactating mammary gland and could also be responsible for the synthesis of complex-type N-linked oligosaccharides in many glycoproteins as well as the carbohydrate moieties of glycolipids.

Cellular Location

[Isoform Long]: Golgi apparatus, Golgi stack membrane; Single-pass type II membrane protein. Cell membrane; Single-pass type II membrane protein Cell surface. Cell projection, filopodium {ECO:0000250|UniProtKB:P15535}. Note=Found in trans cisternae of Golgi but is mainly localized at the plasma membrane (PubMed:1714903) B4GALT1 cell surface expression is regulated by UBE2Q1 (By similarity) {ECO:0000250|UniProtKB:P15535, ECO:0000269|PubMed:1714903} [Processed beta-1,4-galactosyltransferase 1]: Secreted. Note=Soluble form found in body fluids.

Tissue Location


Ubiquitously expressed, but at very low levels in fetal and adult brain

Anti-B4GALT1 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-B4GALT1 Antibody - Images



Western blot analysis of B4GALT1 expression in mouse liver (A), rat liver (B) whole cell lysates.



Immunohistochemical analysis of B4GALT1 staining in human colon cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Anti-B4GALT1 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human B4GALT1. The exact sequence is proprietary.