

## **Anti-GABRG1 Antibody**

Rabbit polyclonal antibody to GABRG1 Catalog # AP60764

## **Specification**

## **Anti-GABRG1 Antibody - Product Information**

Application WB, IHC
Primary Accession Q8N1C3
Other Accession Q9R0Y8
Reactivity Human, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Calculated MW 53595

# **Anti-GABRG1 Antibody - Additional Information**

### **Gene ID 2565**

#### **Other Names**

Gamma-aminobutyric acid receptor subunit gamma-1; GABA(A) receptor subunit gamma-1

### Target/Specificity

Recognizes endogenous levels of GABRG1 protein.

## **Dilution**

WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200) IHC~~1:100~500

#### **Format**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

### Storage

Store at -20 °C. Stable for 12 months from date of receipt

### **Anti-GABRG1 Antibody - Protein Information**

## Name GABRG1 (HGNC:4086)

### **Function**

Gamma subunit of the heteropentameric ligand-gated chloride channel gated by gamma-aminobutyric acid (GABA), a major inhibitory neurotransmitter in the brain (PubMed:<a href="http://www.uniprot.org/citations/10449790" target="\_blank">10449790</a>). GABA-gated chloride channels, also named GABA(A) receptors (GABAAR), consist of five subunits arranged around a central pore and contain GABA active binding site(s) located at the alpha and beta subunit interface(s) (By similarity). When activated by GABA, GABAARs selectively allow the flow of chloride anions across the cell membrane down their electrochemical gradient (PubMed:<a href="http://www.uniprot.org/citations/10449790" target="\_blank">10449790</a>). Chloride



influx into the postsynaptic neuron following GABAAR opening decreases the neuron ability to generate a new action potential, thereby reducing nerve transmission (By similarity).

#### **Cellular Location**


Postsynaptic cell membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein

## **Anti-GABRG1 Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## **Anti-GABRG1 Antibody - Images**



Western blot analysis of GABRG1 expression in HEK293T (A), LOVO (B), Hela (C), mouse brain (D), mouse kidney (E), rat kidney (F) whole cell lysates.



Immunohistochemical analysis of GABRG1 staining in human brain formalin fixed paraffin







embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

# Anti-GABRG1 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human GABRG1. The exact sequence is proprietary.