

**Anti-PORIN Antibody**  
Rabbit polyclonal antibody to PORIN  
Catalog # AP60793**Specification****Anti-PORIN Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P21796</a>
Other Accession	<a href="#">O60932</a>
Reactivity	Human, Mouse, Rat, Rabbit, Monkey, Pig, Bovine, SARS
Host	Rabbit
Clonality	Polyclonal
Calculated MW	30773

**Anti-PORIN Antibody - Additional Information**

Gene ID 7416

**Other Names**

VDAC; Voltage-dependent anion-selective channel protein 1; VDAC-1; hVDAC1; Outer mitochondrial membrane protein porin 1; Plasmalemmal porin; Porin 31HL; Porin 31HM

**Target/Specificity**

Recognizes endogenous levels of PORIN protein.

**Dilution**

WB~~WB (1/500 - 1/1000)

**Format**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**Anti-PORIN Antibody - Protein Information**Name VDAC1 ([HGNC:12669](#))

Synonyms VDAC

**Function**

Non-selective voltage-gated ion channel that mediates the transport of anions and cations through the mitochondrion outer membrane and plasma membrane (PubMed:&lt;a href="http://www.uniprot.org/citations/10661876" target="\_blank"&gt;10661876&lt;/a&gt;, PubMed:&lt;a href="http://www.uniprot.org/citations/11845315" target="\_blank"&gt;11845315&lt;/a&gt;, PubMed:&lt;a href="http://www.uniprot.org/citations/18755977" target="\_blank"&gt;18755977&lt;/a&gt;, PubMed:&lt;a

href="http://www.uniprot.org/citations/30061676" target="\_blank">30061676</a>, PubMed:<a href="http://www.uniprot.org/citations/8420959" target="\_blank">8420959</a>). The channel at the outer mitochondrial membrane allows diffusion of small hydrophilic molecules; in the plasma membrane it is involved in cell volume regulation and apoptosis (PubMed:<a href="http://www.uniprot.org/citations/10661876" target="\_blank">10661876</a>, PubMed:<a href="http://www.uniprot.org/citations/11845315" target="\_blank">11845315</a>, PubMed:<a href="http://www.uniprot.org/citations/18755977" target="\_blank">18755977</a>, PubMed:<a href="http://www.uniprot.org/citations/8420959" target="\_blank">8420959</a>). It adopts an open conformation at low or zero membrane potential and a closed conformation at potentials above 30-40 mV (PubMed:<a href="http://www.uniprot.org/citations/10661876" target="\_blank">10661876</a>, PubMed:<a href="http://www.uniprot.org/citations/18755977" target="\_blank">18755977</a>, PubMed:<a href="http://www.uniprot.org/citations/8420959" target="\_blank">8420959</a>). The open state has a weak anion selectivity whereas the closed state is cation-selective (PubMed:<a href="http://www.uniprot.org/citations/18755977" target="\_blank">18755977</a>, PubMed:<a href="http://www.uniprot.org/citations/8420959" target="\_blank">8420959</a>). Binds various signaling molecules, including the sphingolipid ceramide, the phospholipid phosphatidylcholine, and the sterols cholesterol and oxysterol (PubMed:<a href="http://www.uniprot.org/citations/18755977" target="\_blank">18755977</a>, PubMed:<a href="http://www.uniprot.org/citations/31015432" target="\_blank">31015432</a>). In depolarized mitochondria, acts downstream of PRKN and PINK1 to promote mitophagy or prevent apoptosis; polyubiquitination by PRKN promotes mitophagy, while monoubiquitination by PRKN decreases mitochondrial calcium influx which ultimately inhibits apoptosis (PubMed:<a href="http://www.uniprot.org/citations/32047033" target="\_blank">32047033</a>). May participate in the formation of the permeability transition pore complex (PTPC) responsible for the release of mitochondrial products that triggers apoptosis (PubMed:<a href="http://www.uniprot.org/citations/15033708" target="\_blank">15033708</a>, PubMed:<a href="http://www.uniprot.org/citations/25296756" target="\_blank">25296756</a>). May mediate ATP export from cells (PubMed:<a href="http://www.uniprot.org/citations/30061676" target="\_blank">30061676</a>). Part of a complex composed of HSPA9, ITPR1 and VDAC1 that regulates mitochondrial calcium-dependent apoptosis by facilitating calcium transport from the ER lumen to the mitochondria intermembrane space thus providing calcium for the downstream calcium channel MCU that directly releases it into mitochondria matrix (By similarity). Mediates cytochrome c efflux (PubMed:<a href="http://www.uniprot.org/citations/20230784" target="\_blank">20230784</a>).

#### Cellular Location

Mitochondrion outer membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein. Membrane raft; Multi-pass membrane protein. Note=Found in a complex with HSPA9 and VDAC1 at the endoplasmic reticulum- mitochondria contact sites.  
{ECO:0000250|UniProtKB:Q9Z2L0}

#### Tissue Location

Expressed in erythrocytes (at protein level) (PubMed:27641616). Expressed in heart, liver and skeletal muscle (PubMed:8420959).


#### Anti-PORIN Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-PORIN Antibody - Images



Western blot analysis of PORIN expression in HEK293T (A), Hela (B), mouse brain (C), mouse kidney (D), rat brain (E), rat kidney (F) whole cell lysates.

### Anti-PORIN Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human PORIN. The exact sequence is proprietary.