

**Anti-IRS2 Antibody**  
Rabbit polyclonal antibody to IRS2  
Catalog # AP60877

### Specification

---

#### Anti-IRS2 Antibody - Product Information

Application	WB
Primary Accession	<a href="#">O9Y4H2</a>
Other Accession	<a href="#">P81122</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	137334

#### Anti-IRS2 Antibody - Additional Information

**Gene ID** 8660

#### Other Names

Insulin receptor substrate 2; IRS-2

#### Target/Specificity

Recognizes endogenous levels of IRS2 protein.

#### Dilution

WB~~WB (1/500 - 1/1000)

#### Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

#### Storage

Store at -20 °C. Stable for 12 months from date of receipt

#### Anti-IRS2 Antibody - Protein Information

**Name** IRS2

#### Function

Signaling adapter protein that participates in the signal transduction from two prominent receptor tyrosine kinases, insulin receptor/INSR and insulin-like growth factor I receptor/IGF1R (PubMed: <a href="http://www.uniprot.org/citations/25879670" target="\_blank">25879670</a>). Plays therefore an important role in development, growth, glucose homeostasis as well as lipid metabolism (PubMed: <a href="http://www.uniprot.org/citations/24616100" target="\_blank">24616100</a>). Upon phosphorylation by the insulin receptor, functions as a signaling scaffold that propagates insulin action through binding to SH2 domain-containing proteins including the p85 regulatory subunit of PI3K, NCK1, NCK2, GRB2 or SHP2 (PubMed: <a href="http://www.uniprot.org/citations/15316008" target="\_blank">15316008</a>, PubMed: <a href="http://www.uniprot.org/citations/15316008" target="\_blank">15316008</a>).

<http://www.uniprot.org/citations/19109239> target="\_blank">19109239</a>). Recruitment of GRB2 leads to the activation of the guanine nucleotide exchange factor SOS1 which in turn triggers the Ras/Raf/MEK/MAPK signaling cascade (By similarity). Activation of the PI3K/AKT pathway is responsible for most of insulin metabolic effects in the cell, and the Ras/Raf/MEK/MAPK is involved in the regulation of gene expression and in cooperation with the PI3K pathway regulates cell growth and differentiation. Acts a positive regulator of the Wnt/beta- catenin signaling pathway through suppression of DVL2 autophagy- mediated degradation leading to cell proliferation (PubMed:<a href="http://www.uniprot.org/citations/24616100" target="\_blank">24616100</a>). Plays a role in cell cycle progression by promoting a robust spindle assembly checkpoint (SAC) during M-phase (PubMed:<a href="http://www.uniprot.org/citations/32554797" target="\_blank">32554797</a>). In macrophages, IL4-induced tyrosine phosphorylation of IRS2 leads to the recruitment and activation of phosphoinositide 3-kinase (PI3K) (PubMed:<a href="http://www.uniprot.org/citations/19109239" target="\_blank">19109239</a>).

#### Cellular Location


Cytoplasm, cytosol {ECO:0000250|UniProtKB:P81122}

#### Anti-IRS2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Anti-IRS2 Antibody - Images



Western blot analysis of IRS2 expression in HEK293T (A) whole cell lysates.

#### Anti-IRS2 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human IRS2. The exact sequence is proprietary.