

Anti-iNOS (pY151) Antibody
Rabbit polyclonal antibody to iNOS (pY151)
Catalog # AP60917

Specification

Anti-iNOS (pY151) Antibody - Product Information

Application	WB, E
Primary Accession	P35228
Other Accession	P29477
Reactivity	Human, Mouse, Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	131117

Anti-iNOS (pY151) Antibody - Additional Information

Gene ID 4843

Other Names

NOS2A; Nitric oxide synthase inducible; Hepatocyte NOS; HEP-NOS; Inducible NO synthase; Inducible NOS; iNOS; NOS type II; Peptidyl-cysteine S-nitrosylase NOS2

Target/Specificity

Recognizes endogenous levels of iNOS (pY151) protein.

Dilution

WB~~WB (1/500 - 1/1000), IH (1/50 - 1/100)
E~~WB (1/500 - 1/1000), IH (1/50 - 1/100)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-iNOS (pY151) Antibody - Protein Information

Name NOS2 ([HGNC:7873](#))

Synonyms NOS2A

Function

Produces nitric oxide (NO) which is a messenger molecule with diverse functions throughout the body (PubMed: [7504305](http://www.uniprot.org/citations/7504305), PubMed: [7531687](http://www.uniprot.org/citations/7531687), PubMed: [7544004](http://www.uniprot.org/citations/7544004), PubMed: [7682706](http://www.uniprot.org/citations/7682706))

target="_blank">7682706). In macrophages, NO mediates tumoricidal and bactericidal actions. Also has nitrosylase activity and mediates cysteine S-nitrosylation of cytoplasmic target proteins such PTGS2/COX2 (By similarity). As component of the iNOS-S100A8/9 transnitrosylase complex involved in the selective inflammatory stimulus-dependent S-nitrosylation of GAPDH on 'Cys-247' implicated in regulation of the GAIT complex activity and probably multiple targets including ANXA5, EZR, MSN and VIM (PubMed:25417112). Involved in inflammation, enhances the synthesis of pro-inflammatory mediators such as IL6 and IL8 (PubMed:19688109).

Cellular Location

Cytoplasm, cytosol. Note=Localizes as discrete foci scattered throughout the cytosol and in the presence of SPSB1 and SPSB4, exhibits a more diffuse cytosolic localization.

Tissue Location


Expressed in the liver, retina, bone cells and airway epithelial cells of the lung. Not expressed in the platelets Expressed in chondrocytes (PubMed:7504305)

Anti-iNOS (pY151) Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-iNOS (pY151) Antibody - Images



Western blot analysis of iNOS (pY151) expression in HeLa (A), Jurkat (B), MCF7 (C) whole cell lysates.



Immunohistochemical analysis of iNOS (pY151) staining in human colon cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Direct ELISA antibody dose-response curve using Anti-iNOS (pY151) Antibody. Antigen (phosphopeptide and non-phosphopeptide) concentration is 5 ug/ml. Goat Anti-Rabbit IgG (H&L) - HRP was used as the secondary antibody, and signal was developed by TMB substrate.

Anti-iNOS (pY151) Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human iNOS. The exact sequence is proprietary.