

Anti-ATG4A Antibody
Rabbit polyclonal antibody to ATG4A
Catalog # AP60948

Specification

Anti-ATG4A Antibody - Product Information

Application	WB, IF
Primary Accession	O8WYN0
Other Accession	O8C9S8
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	45378

Anti-ATG4A Antibody - Additional Information

Gene ID 115201

Other Names

APG4A; AUTL2; Cysteine protease ATG4A; AUT-like 2 cysteine endopeptidase; Autophagin-2; Autophagy-related cysteine endopeptidase 2; Autophagy-related protein 4 homolog A; hAPG4A

Target/Specificity

Recognizes endogenous levels of ATG4A protein.

Dilution

WB~~WB (1/500 - 1/1000), IH (1/50 - 1/100), IF/IC (1/100 - 1/500)
IF~~WB (1/500 - 1/1000), IH (1/50 - 1/100), IF/IC (1/100 - 1/500)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-ATG4A Antibody - Protein Information

Name ATG4A {ECO:0000303|Ref.20, ECO:0000312|HGNC:HGNC:16489}

Function

Cysteine protease that plays a key role in autophagy by mediating both proteolytic activation and delipidation of ATG8 family proteins (PubMed:12473658, PubMed:15169837, PubMed:17347651, PubMed:21177865, PubMed:21245471, PubMed:<a

[22302004](http://www.uniprot.org/citations/22302004), PubMed: [32732290](http://www.uniprot.org/citations/32732290)). The protease activity is required for proteolytic activation of ATG8 family proteins: cleaves the C-terminal amino acid of ATG8 proteins to reveal a C-terminal glycine (PubMed: [12473658](http://www.uniprot.org/citations/12473658), PubMed: [15169837](http://www.uniprot.org/citations/15169837), PubMed: [17347651](http://www.uniprot.org/citations/17347651), PubMed: [21177865](http://www.uniprot.org/citations/21177865), PubMed: [21245471](http://www.uniprot.org/citations/21245471), PubMed: [22302004](http://www.uniprot.org/citations/22302004)). Exposure of the glycine at the C-terminus is essential for ATG8 proteins conjugation to phosphatidylethanolamine (PE) and insertion to membranes, which is necessary for autophagy (PubMed: [12473658](http://www.uniprot.org/citations/12473658), PubMed: [15169837](http://www.uniprot.org/citations/15169837), PubMed: [17347651](http://www.uniprot.org/citations/17347651), PubMed: [21177865](http://www.uniprot.org/citations/21177865), PubMed: [21245471](http://www.uniprot.org/citations/21245471), PubMed: [22302004](http://www.uniprot.org/citations/22302004)). Preferred substrate is GABARAP2 followed by MAP1LC3A and GABARAP (PubMed: [12473658](http://www.uniprot.org/citations/12473658), PubMed: [15169837](http://www.uniprot.org/citations/15169837), PubMed: [17347651](http://www.uniprot.org/citations/17347651), PubMed: [21177865](http://www.uniprot.org/citations/21177865), PubMed: [21245471](http://www.uniprot.org/citations/21245471), PubMed: [22302004](http://www.uniprot.org/citations/22302004)). Protease activity is also required to counteract formation of high-molecular weight conjugates of ATG8 proteins (ATG8ylation): acts as a deubiquitinating- like enzyme that removes ATG8 conjugated to other proteins, such as ATG3 (PubMed: [31315929](http://www.uniprot.org/citations/31315929), PubMed: [33773106](http://www.uniprot.org/citations/33773106)). In addition to the protease activity, also mediates delipidation of ATG8 family proteins (PubMed: [29458288](http://www.uniprot.org/citations/29458288), PubMed: [33909989](http://www.uniprot.org/citations/33909989)). Catalyzes delipidation of PE- conjugated forms of ATG8 proteins during macroautophagy (PubMed: [29458288](http://www.uniprot.org/citations/29458288), PubMed: [33909989](http://www.uniprot.org/citations/33909989)). Compared to ATG4B, the major protein for proteolytic activation of ATG8 proteins, shows weaker ability to cleave the C-terminal amino acid of ATG8 proteins, while it displays stronger delipidation activity (PubMed: [29458288](http://www.uniprot.org/citations/29458288)). Involved in phagophore growth during mitophagy independently of its protease activity and of ATG8 proteins: acts by regulating ATG9A trafficking to mitochondria and promoting phagophore-endoplasmic reticulum contacts during the lipid transfer phase of mitophagy (PubMed: [33773106](http://www.uniprot.org/citations/33773106)).

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:Q8BGE6}.


Anti-ATG4A Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-ATG4A Antibody - Images



Western blot analysis of ATG4A expression in HCT116 (A), HepG2 (B), CT26 (C), PC12 (D) whole cell lysates.



Immunohistochemical analysis of ATG4A staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of ATG4A staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with Alexa Fluor 647-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Anti-ATG4A Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human ATG4A. The exact sequence is proprietary.