

Anti-WEE1 (pS642) Antibody
Rabbit polyclonal antibody to WEE1 (pS642)
Catalog # AP61084

Specification

Anti-WEE1 (pS642) Antibody - Product Information

| | |
|-------------------|------------------------|
| Application | WB |
| Primary Accession | P30291 |
| Other Accession | P47810 |
| Reactivity | Human, Mouse, Rat |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 71597 |

Anti-WEE1 (pS642) Antibody - Additional Information

Gene ID 7465

Other Names

Wee1-like protein kinase; WEE1hu; Wee1A kinase

Target/Specificity

Recognizes endogenous levels of WEE1 (pS642) protein.

Dilution

WB~~WB (1/500 - 1/1000)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-WEE1 (pS642) Antibody - Protein Information

Name WEE1 {ECO:0000303|PubMed:8348613, ECO:0000312|HGNC:HGNC:12761}

Function

Acts as a negative regulator of entry into mitosis (G2 to M transition) by protecting the nucleus from cytoplasmically activated cyclin B1-complexed CDK1 before the onset of mitosis by mediating phosphorylation of CDK1 on 'Tyr-15' (PubMed:15070733, PubMed:7743995, PubMed:8348613, PubMed:8428596). Specifically phosphorylates and inactivates cyclin B1-complexed CDK1 reaching a maximum during G2 phase and a minimum as cells enter M phase (PubMed:7743995, PubMed:8348613, PubMed:8428596). Phosphorylation of cyclin B1-CDK1 occurs exclusively on 'Tyr-15' and phosphorylation of monomeric CDK1 does not occur (PubMed:7743995, PubMed:8348613, PubMed:8428596). Its activity increases during S and G2 phases and decreases at M phase when it is hyperphosphorylated (PubMed:7743995). A correlated decrease in protein level occurs at M/G1 phase, probably due to its degradation (PubMed:7743995).

Cellular Location


Nucleus.

Anti-WEE1 (pS642) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-WEE1 (pS642) Antibody - Images



Western blot analysis of WEE1 (pS642) expression in mouse brain (A), rat brain (B) whole cell lysates.

Anti-WEE1 (pS642) Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human WEE1. The exact sequence is proprietary.