

## Anti-Histone H100 (AcK163) Antibody

Rabbit polyclonal antibody to Histone H1oo (AcK163) Catalog # AP61134

#### **Specification**

## Anti-Histone H100 (AcK163) Antibody - Product Information

Application WB
Primary Accession Q8IZA3
Other Accession Q8VIK3

Reactivity Human, Mouse, Rat Host Rabbit

Clonality Polyclonal Calculated MW 35813

# Anti-Histone H1oo (AcK163) Antibody - Additional Information

**Gene ID 132243** 

#### **Other Names**

H100; OSH1; Histone H100; Oocyte-specific histone H1; Oocyte-specific linker histone H1; osH1

#### Target/Specificity

Recognizes endogenous levels of Histone H100 (AcK163) protein.

## **Dilution**

WB~~WB (1/500 - 1/1000), IH (1/50 - 1/200)

#### **Format**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

## **Storage**

Store at -20 °C. Stable for 12 months from date of receipt

# Anti-Histone H100 (AcK163) Antibody - Protein Information

Name H1-8 (<u>HGNC:18463</u>)

#### **Function**

May play a key role in the control of gene expression during oogenesis and early embryogenesis, presumably through the perturbation of chromatin structure. Essential for meiotic maturation of germinal vesicle-stage oocytes. The somatic type linker histone H1c is rapidly replaced by H1oo in a donor nucleus transplanted into an oocyte. The greater mobility of H1oo as compared to H1c may contribute to this rapid replacement and increased instability of the embryonic chromatin structure. The rapid replacement of H1c with H1oo may play an important role in nuclear remodeling (By similarity).

## **Cellular Location**



Cytoplasm. Nucleus {ECO:0000255|PROSITE-ProRule:PRU00837}. Chromosome {ECO:0000255|PROSITE-ProRule:PRU00837}


**Tissue Location** Oocyte-specific..

# Anti-Histone H1oo (AcK163) Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# Anti-Histone H100 (AcK163) Antibody - Images



Western blot analysis of Histone H1oo (AcK163) expression in A2780 (A), U87MG (B), mouse testis (C), rat brain (D) whole cell lysates.



Immunohistochemical analysis of Histone H100 (AcK163) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated





Tel: 858.875.1900 Fax: 858.875.1999

antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

# Anti-Histone H1oo (AcK163) Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Histone H100 (AcK163). The exact sequence is proprietary.