

Anti-NDUFC2 Antibody
Rabbit polyclonal antibody to NDUFC2
Catalog # AP61194

Specification

Anti-NDUFC2 Antibody - Product Information

Application	WB
Primary Accession	O95298
Other Accession	O9C054
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	14188

Anti-NDUFC2 Antibody - Additional Information

Gene ID 4718

Other Names

NADH dehydrogenase [ubiquinone] 1 subunit C2; Complex I-B14.5b; CI-B14.5b; Human lung cancer oncogene 1 protein; HLC-1; NADH-ubiquinone oxidoreductase subunit B14.5b

Target/Specificity

Recognizes endogenous levels of NDUFC2 protein.

Dilution

WB~~WB (1/500 - 1/1000), IH (1/50 - 1/200)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-NDUFC2 Antibody - Protein Information

Name NDUFC2 ([HGNC:7706](#))

Function

Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed not to be involved in catalysis but required for the complex assembly. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.

Cellular Location


Mitochondrion inner membrane; Single-pass membrane protein; Matrix side

Anti-NDUFC2 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-NDUFC2 Antibody - Images



Western blot analysis of NDUFC2 expression in A375 (A), PC3 (B), H9C2 (C), AML12 (D) whole cell lysates.



Immunohistochemical analysis of NDUFC2 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Anti-NDUFC2 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human NDUFC2. The exact sequence is proprietary.