

Anti-Frizzled 1 Antibody
Rabbit polyclonal antibody to Frizzled 1
Catalog # AP61230**Specification**

Anti-Frizzled 1 Antibody - Product Information

Application	WB, IF
Primary Accession	O9UP38
Other Accession	O70421
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	71158

Anti-Frizzled 1 Antibody - Additional Information**Gene ID** 8321**Other Names**

Frizzled-1; Fz-1; hFz1; FzE1

Target/Specificity

Recognizes endogenous levels of Frizzled 1 protein.

Dilution

WB~~WB (1/500 - 1/1000), IF/IC (1/100 - 1/500)

IF~~WB (1/500 - 1/1000), IF/IC (1/100 - 1/500)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-Frizzled 1 Antibody - Protein Information**Name** FZD1**Function**

Receptor for Wnt proteins (PubMed:10557084). Activated by WNT3A, WNT3, WNT1 and to a lesser extent WNT2, but apparently not by WNT4, WNT5A, WNT5B, WNT6, WNT7A or WNT7B (PubMed:10557084). Contradictory results showing activation by WNT7B have been described for mouse (By similarity). Functions in the canonical Wnt/beta-catenin signaling pathway (PubMed:10557084). The canonical Wnt/beta-catenin signaling pathway leads to the activation of disheveled proteins,

inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin and activation of Wnt target genes (PubMed:10557084). A second signaling pathway involving PKC and calcium fluxes has been seen for some family members, but it is not yet clear if it represents a distinct pathway or if it can be integrated in the canonical pathway, as PKC seems to be required for Wnt-mediated inactivation of GSK-3 kinase. Both pathways seem to involve interactions with G-proteins. May be involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and/or in differentiated tissues (Probable).

Cellular Location

Cell membrane; Multi-pass membrane protein


Tissue Location

Expressed in adult heart, placenta, lung, kidney, pancreas, prostate, and ovary and in fetal lung and kidney


Anti-Frizzled 1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Frizzled 1 Antibody - Images

Western blot analysis of Frizzled 1 expression in A549 (A), Panc1 (B), CT26 (C) whole cell lysates.



Immunofluorescent analysis of Frizzled 1 staining in HuvEc cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a Alexa Fluor 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Anti-Frizzled 1 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human Frizzled 1. The exact sequence is proprietary.