

Anti-Glucocorticoid Receptor (pS203) Antibody
Rabbit polyclonal antibody to Glucocorticoid Receptor (pS203)
Catalog # AP61259

Specification

Anti-Glucocorticoid Receptor (pS203) Antibody - Product Information

Application	WB
Primary Accession	P04150
Other Accession	P06537
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	85659

Anti-Glucocorticoid Receptor (pS203) Antibody - Additional Information

Gene ID 2908

Other Names

GRL; Glucocorticoid receptor; GR; Nuclear receptor subfamily 3 group C member 1

Target/Specificity

Recognizes endogenous levels of Glucocorticoid Receptor (pS203) protein.

Dilution

WB~~WB (1/500 - 1/1000)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-Glucocorticoid Receptor (pS203) Antibody - Protein Information

Name NR3C1 ([HGNC:7978](#))

Synonyms GRL

Function

Receptor for glucocorticoids (GC) (PubMed: [27120390](http://www.uniprot.org/citations/27120390), PubMed: [37478846](http://www.uniprot.org/citations/37478846)). Has a dual mode of action: as a transcription factor that binds to glucocorticoid response elements (GRE), both for nuclear and mitochondrial DNA, and as a modulator of other transcription factors (PubMed: [28139699](http://www.uniprot.org/citations/28139699)). Affects inflammatory responses, cellular proliferation and differentiation in target tissues. Involved in

chromatin remodeling (PubMed:9590696). Plays a role in rapid mRNA degradation by binding to the 5' UTR of target mRNAs and interacting with PNRC2 in a ligand-dependent manner which recruits the RNA helicase UPF1 and the mRNA-decapping enzyme DCP1A, leading to RNA decay (PubMed:25775514). Could act as a coactivator for STAT5-dependent transcription upon growth hormone (GH) stimulation and could reveal an essential role of hepatic GR in the control of body growth (By similarity).

Cellular Location

[Isoform Alpha]: Cytoplasm. Nucleus. Mitochondrion. Cytoplasm, cytoskeleton, spindle. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Note=After ligand activation, translocates from the cytoplasm to the nucleus. In the presence of NR1D1 shows a time-dependent subcellular localization, localizing to the cytoplasm at ZT8 and to the nucleus at ZT20 (By similarity). Lacks this diurnal pattern of localization in the absence of NR1D1, localizing to both nucleus and the cytoplasm at ZT8 and ZT20 (By similarity).

{ECO:0000250|UniProtKB:P06537, ECO:0000269|PubMed:18838540,

ECO:0000269|PubMed:27120390, ECO:0000269|PubMed:8621628} [Isoform Alpha-B]: Nucleus.

Cytoplasm Note=After ligand activation, translocates from the cytoplasm to the nucleus.

Tissue Location


Widely expressed including bone, stomach, lung, liver, colon, breast, ovary, pancreas and kidney (PubMed:25847991). In the heart, detected in left and right atria, left and right ventricles, aorta, apex, intraventricular septum, and atrioventricular node as well as whole adult and fetal heart (PubMed:10902803) [Isoform Alpha-2]: Widely expressed.

Anti-Glucocorticoid Receptor (pS203) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Glucocorticoid Receptor (pS203) Antibody - Images



Western blot analysis of Glucocorticoid Receptor (pS203) expression in Hela (A), A375 (B), LO2 (C), AML12 (D), PMVEC (E) whole cell lysates.

Anti-Glucocorticoid Receptor (pS203) Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Glucocorticoid Receptor (pS203). The exact sequence is proprietary.