

**Anti-NFAT3 (pS168/S170) Antibody**  
**Rabbit polyclonal antibody to NFAT3 (pS168/S170)**  
**Catalog # AP61272**

## Specification

---

### Anti-NFAT3 (pS168/S170) Antibody - Product Information

Application	WB
Primary Accession	<a href="#">O14934</a>
Other Accession	<a href="#">O8K120</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	95449

### Anti-NFAT3 (pS168/S170) Antibody - Additional Information

**Gene ID** 4776

#### Other Names

NFAT3; Nuclear factor of activated T-cells cytoplasmic 4; NF-ATc4; NFATc4; T-cell transcription factor NFAT3; NF-AT3

#### Target/Specificity

Recognizes endogenous levels of NFAT3 (pS168/S170) protein.

#### Dilution

WB~~WB (1/500 - 1/1000), IH (1/50 - 1/200)

#### Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

#### Storage

Store at -20 °C. Stable for 12 months from date of receipt

### Anti-NFAT3 (pS168/S170) Antibody - Protein Information

**Name** NFATC4

#### Function

Ca(2+)-regulated transcription factor that is involved in several processes, including the development and function of the immune, cardiovascular, musculoskeletal, and nervous systems (PubMed: [11514544](http://www.uniprot.org/citations/11514544)), PubMed: [11997522](http://www.uniprot.org/citations/11997522), PubMed: [17213202](http://www.uniprot.org/citations/17213202), PubMed: [17875713](http://www.uniprot.org/citations/17875713), PubMed: [18668201](http://www.uniprot.org/citations/18668201), PubMed: [25663301](http://www.uniprot.org/citations/25663301)),

PubMed: [7749981](http://www.uniprot.org/citations/7749981)). Involved in T-cell activation, stimulating the transcription of cytokine genes, including that of IL2 and IL4 (PubMed: [18347059](http://www.uniprot.org/citations/18347059)), PubMed: [18668201](http://www.uniprot.org/citations/18668201), PubMed: [7749981](http://www.uniprot.org/citations/7749981)). Along with NFATC3, involved in embryonic heart development. Following JAK/STAT signaling activation and as part of a complex with NFATC3 and STAT3, binds to the alpha-beta E4 promoter region of CRYAB and activates transcription in cardiomyocytes (By similarity). Involved in mitochondrial energy metabolism required for cardiac morphogenesis and function (By similarity). Transactivates many genes involved in the cardiovascular system, including AGTR2, NPPB/BNP (in synergy with GATA4), NPPA/ANP/ANF and MYH7/beta-MHC (By similarity). Involved in the regulation of adult hippocampal neurogenesis. Involved in BDNF-driven pro-survival signaling in hippocampal adult-born neurons. Involved in the formation of long-term spatial memory and long-term potentiation (By similarity). In cochlear nucleus neurons, may play a role in deafferentation-induced apoptosis during the developmental critical period, when auditory neurons depend on afferent input for survival (By similarity). Binds to and activates the BACE1/Beta-secretase 1 promoter, hence may regulate the proteolytic processing of the amyloid precursor protein (APP) (PubMed: [25663301](http://www.uniprot.org/citations/25663301)). Plays a role in adipocyte differentiation (PubMed: [11997522](http://www.uniprot.org/citations/11997522)). May be involved in myoblast differentiation into myotubes (PubMed: [17213202](http://www.uniprot.org/citations/17213202)). Binds the consensus DNA sequence 5'-GGAAAAT-3' (Probable). In the presence of CREBBP, activates TNF transcription (PubMed: [11514544](http://www.uniprot.org/citations/11514544)). Binds to PPARG gene promoter and regulates its activity (PubMed: [11997522](http://www.uniprot.org/citations/11997522)). Binds to PPARG and REG3G gene promoters (By similarity).

#### Cellular Location

Cytoplasm. Nucleus. Note=When hyperphosphorylated, localizes in the cytosol. When intracellular Ca(2+) levels increase, dephosphorylation by calcineurin/PPP3CA leads to translocation into the nucleus (PubMed:11997522, PubMed:18347059). MAPK7/ERK5 and MTOR regulate NFATC4 nuclear export through phosphorylation at Ser-168 and Ser-170 (PubMed:18347059).

#### Tissue Location


Widely expressed, with high levels in placenta, lung, kidney, testis and ovary (PubMed:18675896). Weakly expressed in spleen and thymus (PubMed:18675896). In the hippocampus, expressed in the granular layer of the dentate gyrus, in the pyramidal neurons of CA3 region, and in the hippocampal fissure (PubMed:18675896). Expressed in the heart (at protein level) (PubMed:12370307)

### Anti-NFAT3 (pS168/S170) Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-NFAT3 (pS168/S170) Antibody - Images



Western blot analysis of NFAT3 (pS168/S170) expression in A375 (A), HEK293T (B), CT26 (C) whole cell lysates.



Immunohistochemical analysis of NFAT3 (pS168/S170) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

#### **Anti-NFAT3 (pS168/S170) Antibody - Background**

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human NFAT3 (pS168/S170). The exact sequence is proprietary.