

**Anti-FHIT (pY114) Antibody**  
Rabbit polyclonal antibody to FHIT (pY114)  
Catalog # AP61277

## Specification

---

### Anti-FHIT (pY114) Antibody - Product Information

Application	WB
Primary Accession	<a href="#">P49789</a>
Other Accession	<a href="#">O89106</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	16858

### Anti-FHIT (pY114) Antibody - Additional Information

Gene ID 2272

#### Other Names

Bis(5'-adenosyl)-triphosphatase; AP3A hydrolase; AP3Aase; Diadenosine 5'5'''-P1P3-triphosphate hydrolase; Dinucleosidetriphosphatase; Fragile histidine triad protein

#### Target/Specificity

Recognizes endogenous levels of FHIT (pY114) protein.

#### Dilution

WB~~WB (1/500 - 1/1000), IH (1/50 - 1/200)

#### Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

#### Storage

Store at -20 °C. Stable for 12 months from date of receipt

### Anti-FHIT (pY114) Antibody - Protein Information

Name FHIT

#### Function

Possesses dinucleoside triphosphate hydrolase activity (PubMed:<a href="http://www.uniprot.org/citations/12574506" target="\_blank">12574506</a>, PubMed:<a href="http://www.uniprot.org/citations/15182206" target="\_blank">15182206</a>, PubMed:<a href="http://www.uniprot.org/citations/8794732" target="\_blank">8794732</a>, PubMed:<a href="http://www.uniprot.org/citations/9323207" target="\_blank">9323207</a>, PubMed:<a href="http://www.uniprot.org/citations/9543008" target="\_blank">9543008</a>, PubMed:<a href="http://www.uniprot.org/citations/9576908" target="\_blank">9576908</a>). Cleaves P(1)-P(3)-bis(5'-adenosyl) triphosphate (Ap3A) to yield AMP and ADP (PubMed:<a

href="http://www.uniprot.org/citations/12574506" target="\_blank">12574506</a>, PubMed:<a href="http://www.uniprot.org/citations/15182206" target="\_blank">15182206</a>, PubMed:<a href="http://www.uniprot.org/citations/8794732" target="\_blank">8794732</a>, PubMed:<a href="http://www.uniprot.org/citations/9323207" target="\_blank">9323207</a>, PubMed:<a href="http://www.uniprot.org/citations/9543008" target="\_blank">9543008</a>, PubMed:<a href="http://www.uniprot.org/citations/9576908" target="\_blank">9576908</a>). Can also hydrolyze P(1)-P(4)-bis(5'-adenosyl) tetraphosphate (Ap4A), but has extremely low activity with ATP (PubMed:<a href="http://www.uniprot.org/citations/8794732" target="\_blank">8794732</a>). Exhibits adenylylsulfatase activity, hydrolyzing adenosine 5'-phosphosulfate to yield AMP and sulfate (PubMed:<a href="http://www.uniprot.org/citations/18694747" target="\_blank">18694747</a>). Exhibits adenosine 5'-monophosphoramidase activity, hydrolyzing purine nucleotide phosphoramidates with a single phosphate group such as adenosine 5'-monophosphoramidate (AMP-NH<sub>2</sub>) to yield AMP and NH<sub>2</sub> (PubMed:<a href="http://www.uniprot.org/citations/18694747" target="\_blank">18694747</a>). Exhibits adenylylsulfate-ammonia adenylyltransferase, catalyzing the ammonolysis of adenosine 5'-phosphosulfate resulting in the formation of adenosine 5'-phosphoramidate (PubMed:<a href="http://www.uniprot.org/citations/26181368" target="\_blank">26181368</a>). Also catalyzes the ammonolysis of adenosine 5-phosphorofluoridate and diadenosine triphosphate (PubMed:<a href="http://www.uniprot.org/citations/26181368" target="\_blank">26181368</a>). Modulates transcriptional activation by CTNBN1 and thereby contributes to regulate the expression of genes essential for cell proliferation and survival, such as CCND1 and BIRC5 (PubMed:<a href="http://www.uniprot.org/citations/18077326" target="\_blank">18077326</a>). Plays a role in the induction of apoptosis via SRC and AKT1 signaling pathways (PubMed:<a href="http://www.uniprot.org/citations/16407838" target="\_blank">16407838</a>). Inhibits MDM2-mediated proteasomal degradation of p53/TP53 and thereby plays a role in p53/TP53-mediated apoptosis (PubMed:<a href="http://www.uniprot.org/citations/15313915" target="\_blank">15313915</a>). Induction of apoptosis depends on the ability of FHIT to bind P(1)-P(3)-bis(5'-adenosyl) triphosphate or related compounds, but does not require its catalytic activity, it may in part come from the mitochondrial form, which sensitizes the low-affinity Ca(2+) transporters, enhancing mitochondrial calcium uptake (PubMed:<a href="http://www.uniprot.org/citations/12574506" target="\_blank">12574506</a>, PubMed:<a href="http://www.uniprot.org/citations/19622739" target="\_blank">19622739</a>). Functions as a tumor suppressor (By similarity).

#### Cellular Location

Cytoplasm. Mitochondrion. Nucleus

#### Tissue Location


Low levels expressed in all tissues tested. Phospho-FHIT observed in liver and kidney, but not in brain and lung Phospho-FHIT undetected in all tested human tumor cell lines

#### Anti-FHIT (pY114) Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Anti-FHIT (pY114) Antibody - Images



Western blot analysis of FHIT (pY114) expression in NIH3T3 (A), U87MG (B), Beas2B (C), SGC7901 (D) whole cell lysates.



Immunohistochemical analysis of FHIT (pY114) staining in human colon cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

#### **Anti-FHIT (pY114) Antibody - Background**

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human FHIT (pY114). The exact sequence is proprietary.