

**Anti-Granulin Antibody**  
Rabbit polyclonal antibody to Granulin  
Catalog # AP61369

### Specification

---

#### Anti-Granulin Antibody - Product Information

Application	<b>WB</b>
Primary Accession	<a href="#">P28799</a>
Reactivity	<b>Human, Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>63544</b>

#### Anti-Granulin Antibody - Additional Information

**Gene ID** 2896

#### Other Names

Granulins; Proepithelin; PEPI

#### Target/Specificity

Recognizes endogenous levels of Granulin protein.

#### Dilution

WB~~WB (1/500 - 1/1000)

#### Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

#### Storage

Store at -20 °C. Stable for 12 months from date of receipt

#### Anti-Granulin Antibody - Protein Information

**Name** GRN ([HGNC:4601](#))

#### Function

Secreted protein that acts as a key regulator of lysosomal function and as a growth factor involved in inflammation, wound healing and cell proliferation (PubMed: [12526812](http://www.uniprot.org/citations/12526812), PubMed: [18378771](http://www.uniprot.org/citations/18378771), PubMed: [28073925](http://www.uniprot.org/citations/28073925), PubMed: [28453791](http://www.uniprot.org/citations/28453791), PubMed: [28541286](http://www.uniprot.org/citations/28541286)). Regulates protein trafficking to lysosomes and, also the activity of lysosomal enzymes (PubMed: [28453791](http://www.uniprot.org/citations/28453791), PubMed: [28541286](http://www.uniprot.org/citations/28541286)). Facilitates

also the acidification of lysosomes, causing degradation of mature CTSD by CTSB (PubMed:<a href="http://www.uniprot.org/citations/28073925" target="\_blank">28073925</a>). In addition, functions as a wound-related growth factor that acts directly on dermal fibroblasts and endothelial cells to promote division, migration and the formation of capillary-like tubule structures (By similarity). Also promotes epithelial cell proliferation by blocking TNF-mediated neutrophil activation preventing release of oxidants and proteases (PubMed:<a href="http://www.uniprot.org/citations/12526812" target="\_blank">12526812</a>). Moreover, modulates inflammation in neurons by preserving neurons survival, axonal outgrowth and neuronal integrity (PubMed:<a href="http://www.uniprot.org/citations/18378771" target="\_blank">18378771</a>).

### Cellular Location

Secreted. Lysosome Note=Endocytosed by SORT1 and delivered to lysosomes (PubMed:21092856, PubMed:28073925). Targeted to lysosome by PSAP via M6PR and LRP1, in both biosynthetic and endocytic pathways (PubMed:26370502, PubMed:28073925). Co-localized with GBA1 in the intracellular trafficking compartments until to lysosome (By similarity) {ECO:0000250|UniProtKB:P28798, ECO:0000269|PubMed:21092856, ECO:0000269|PubMed:26370502, ECO:0000269|PubMed:28073925}

### Tissue Location


In myelogenous leukemic cell lines of promonocytic, promyelocytic, and proerythroid lineage, in fibroblasts, and very strongly in epithelial cell lines. Present in inflammatory cells and bone marrow. Highest levels in kidney

### Anti-Granulin Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-Granulin Antibody - Images



Western blot analysis of Granulin expression in rat kidney (A), HEK293T (B), Jurkat (C), Beas2B

(D) whole cell lysates.

### **Anti-Granulin Antibody - Background**

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Granulin. The exact sequence is proprietary.