

Anti-CD144 Antibody
Rabbit polyclonal antibody to CD144
Catalog # AP61383

Specification

Anti-CD144 Antibody - Product Information

Application	WB
Primary Accession	P33151
Other Accession	P55284
Reactivity	Human, Mouse, Chicken, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	87528

Anti-CD144 Antibody - Additional Information

Gene ID 1003

Other Names

Cadherin-5; 7B4 antigen; Vascular endothelial cadherin; VE-cadherin; CD144

Target/Specificity

Recognizes endogenous levels of CD144 protein.

Dilution

WB~~WB (1/500 - 1/1000), IH (1/50 - 1/200)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-CD144 Antibody - Protein Information

Name CDH5 ([HGNC:1764](#))

Function

Cadherins are calcium-dependent cell adhesion proteins (By similarity). They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types (PubMed:21269602). This cadherin may play an important role in endothelial cell biology through control of the cohesion and organization of the intercellular junctions (By similarity). It associates with alpha-catenin forming a link to the cytoskeleton (PubMed:10861224). Plays a role in coupling actin fibers to cell junctions in endothelial cells, via acting as a cell junctional complex anchor for AMOTL2 and MAGI1 (By

similarity). Acts in concert with KRIT1 and PALS1 to establish and maintain correct endothelial cell polarity and vascular lumen (By similarity). These effects are mediated by recruitment and activation of the Par polarity complex and RAP1B (PubMed:20332120). Required for activation of PRKCZ and for the localization of phosphorylated PRKCZ, PARD3, TIAM1 and RAP1B to the cell junction (PubMed:20332120).

Cellular Location

Cell junction, adherens junction. Cell membrane; Single-pass type I membrane protein Cytoplasm {ECO:0000250|UniProtKB:P55284}. Note=Found at cell-cell boundaries and probably at cell-matrix boundaries. KRIT1 and CDH5 reciprocally regulate their localization to endothelial cell-cell junctions.

Tissue Location


Endothelial tissues and brain.

Anti-CD144 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CD144 Antibody - Images



Western blot analysis of CD144 expression in A375 (A), U87MG (B), mouse brain (C), rat brain (D) whole cell lysates.



Immunohistochemical analysis of CD144 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Anti-CD144 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human CD144. The exact sequence is proprietary.