

**Anti-TRAF6 Antibody**  
Rabbit polyclonal antibody to TRAF6  
Catalog # AP61519

### Specification

---

#### Anti-TRAF6 Antibody - Product Information

Application	WB
Primary Accession	<a href="#">O9Y4K3</a>
Other Accession	<a href="#">P70196</a>
Reactivity	Human, Mouse, Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	59573

#### Anti-TRAF6 Antibody - Additional Information

Gene ID 7189

#### Other Names

RNF85; TNF receptor-associated factor 6; E3 ubiquitin-protein ligase TRAF6; Interleukin-1 signal transducer; RING finger protein 85

#### Target/Specificity

Recognizes endogenous levels of TRAF6 protein.

#### Dilution

WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200)

#### Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

#### Storage

Store at -20 °C. Stable for 12 months from date of receipt

#### Anti-TRAF6 Antibody - Protein Information

Name TRAF6

Synonyms RNF85

#### Function

E3 ubiquitin ligase that, together with UBE2N and UBE2V1, mediates the synthesis of 'Lys-63'-linked-polyubiquitin chains conjugated to proteins, such as ECSIT, IKBKG, IRAK1, AKT1 and AKT2 (PubMed: [11057907](http://www.uniprot.org/citations/11057907) target="\_blank">11057907</a>, PubMed: [18347055](http://www.uniprot.org/citations/18347055) target="\_blank">18347055</a>, PubMed: [19465916](http://www.uniprot.org/citations/19465916) target="\_blank">19465916</a>, PubMed: [19713527](http://www.uniprot.org/citations/19713527) target="\_blank">19713527</a>)

target="\_blank">19713527</a>, PubMed:<a href="http://www.uniprot.org/citations/31620128" target="\_blank">31620128</a>). Also mediates ubiquitination of free/unanchored polyubiquitin chain that leads to MAP3K7 activation (PubMed:<a href="http://www.uniprot.org/citations/19675569" target="\_blank">19675569</a>). Leads to the activation of NF-kappa-B and JUN (PubMed:<a href="http://www.uniprot.org/citations/16378096" target="\_blank">16378096</a>, PubMed:<a href="http://www.uniprot.org/citations/17135271" target="\_blank">17135271</a>, PubMed:<a href="http://www.uniprot.org/citations/17703191" target="\_blank">17703191</a>). Seems to also play a role in dendritic cells (DCs) maturation and/or activation (By similarity). Represses c-Myb-mediated transactivation, in B-lymphocytes (PubMed:<a href="http://www.uniprot.org/citations/18093978" target="\_blank">18093978</a>, PubMed:<a href="http://www.uniprot.org/citations/18758450" target="\_blank">18758450</a>). Adapter protein that seems to play a role in signal transduction initiated via TNF receptor, IL-1 receptor and IL-17 receptor (PubMed:<a href="http://www.uniprot.org/citations/12140561" target="\_blank">12140561</a>, PubMed:<a href="http://www.uniprot.org/citations/19825828" target="\_blank">19825828</a>, PubMed:<a href="http://www.uniprot.org/citations/8837778" target="\_blank">8837778</a>). Regulates osteoclast differentiation by mediating the activation of adapter protein complex 1 (AP-1) and NF-kappa-B, in response to RANK-L stimulation (By similarity). Together with MAP3K8, mediates CD40 signals that activate ERK in B-cells and macrophages, and thus may play a role in the regulation of immunoglobulin production (By similarity). Participates also in the TCR signaling by ubiquitinating LAT (PubMed:<a href="http://www.uniprot.org/citations/23514740" target="\_blank">23514740</a>, PubMed:<a href="http://www.uniprot.org/citations/25907557" target="\_blank">25907557</a>).

#### Cellular Location

Cytoplasm. Cytoplasm, cell cortex. Nucleus. Lipid droplet {ECO:0000250|UniProtKB:P70196}. Note=Found in the nuclei of some aggressive B-cell lymphoma cell lines as well as in the nuclei of both resting and activated T- and B-lymphocytes. Found in punctate nuclear body protein complexes. Ubiquitination may occur in the cytoplasm and sumoylation in the nucleus. RSAD2/viperin recruits it to the lipid droplet (By similarity).

#### Tissue Location

Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas


#### Anti-TRAF6 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Anti-TRAF6 Antibody - Images





Western blot analysis of TRAF6 expression in Hela (A), K562 (B), Rat kidney (C), Rat heart (D), mouse kidney (E) whole cell lysates.



Immunohistochemical analysis of TRAF6 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

#### **Anti-TRAF6 Antibody - Background**

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human TRAF6. The exact sequence is proprietary.