

Anti-Peroxiredoxin 1 Antibody
Mouse monoclonal antibody to Peroxiredoxin 1
Catalog # AP61594**Specification**

Anti-Peroxiredoxin 1 Antibody - Product Information

| | |
|-------------------|--------------------------|
| Application | WB, IF |
| Primary Accession | Q06830 |
| Other Accession | P35700 |
| Reactivity | Human, Mouse, Rat |
| Host | Mouse |
| Clonality | Monoclonal |
| Calculated MW | 22110 |

Anti-Peroxiredoxin 1 Antibody - Additional Information**Gene ID** 5052**Other Names**

PAGA; PAGB; TDPX2; Peroxiredoxin-1; Natural killer cell-enhancing factor A; NKEF-A; Proliferation-associated gene protein; PAG; Thioredoxin peroxidase 2; Thioredoxin-dependent peroxide reductase 2

Target/Specificity

Recognizes endogenous levels of Peroxiredoxin 1 protein.

Dilution

WB~~WB (1/1000 - 1/3000), IF/IC (1/100 - 1/200)
IF~~WB (1/1000 - 1/3000), IF/IC (1/100 - 1/200)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-Peroxiredoxin 1 Antibody - Protein Information**Name** PRDX1**Synonyms** PAGA, PAGB, TDPX2**Function**

Thiol-specific peroxidase that catalyzes the reduction of hydrogen peroxide and organic hydroperoxides to water and alcohols, respectively. Plays a role in cell protection against oxidative stress by detoxifying peroxides and as sensor of hydrogen peroxide-mediated signaling events. Might participate in the signaling cascades of growth factors and tumor necrosis factor-alpha by

regulating the intracellular concentrations of H₂O₂ (PubMed:9497357). Reduces an intramolecular disulfide bond in GDPD5 that gates the ability to GDPD5 to drive postmitotic motor neuron differentiation (By similarity).

Cellular Location


Cytoplasm. Melanosome Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV

Anti-Peroxiredoxin 1 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Peroxiredoxin 1 Antibody - Images



Western blot analysis of Peroxiredoxin 1 expression in MCF7 (A), mouse brain (B), rat kidney (C) whole cell lysates.



Immunofluorescent analysis of Peroxiredoxin 1 staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a FITC-conjugated secondary antibody (green) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Anti-Peroxiredoxin 1 Antibody - Background

Recombinant protein corresponding to human Peroxiredoxin 1.