

**Anti-LC3B Antibody**  
Rabbit polyclonal antibody to LC3B  
Catalog # AP61601**Specification**

---

**Anti-LC3B Antibody - Product Information**

Application	WB, IF
Primary Accession	<a href="#">O9GZQ8</a>
Other Accession	<a href="#">O9CQV6</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	14688

**Anti-LC3B Antibody - Additional Information**

Gene ID 81631

**Other Names**

MAP1ALC3; Microtubule-associated proteins 1A/1B light chain 3B; Autophagy-related protein LC3 B; Autophagy-related ubiquitin-like modifier LC3 B; MAP1 light chain 3-like protein 2; MAP1A/MAP1B light chain 3 B; MAP1A/MAP1B LC3 B; Microtubule-associated protein 1 light chain 3 beta

**Target/Specificity**

Recognizes endogenous levels of LC3B protein.

**Dilution**

WB~~WB (1/1000 - 1/2000), IH (1/200 - 1/500), IF/IC (1/100 - 1/200)  
IF~~WB (1/1000 - 1/2000), IH (1/200 - 1/500), IF/IC (1/100 - 1/200)

**Format**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**Anti-LC3B Antibody - Protein Information**

Name MAP1LC3B ([HGNC:13352](#))

Synonyms MAP1ALC3

**Function**

Ubiquitin-like modifier involved in formation of autophagosomal vacuoles (autophagosomes) (PubMed: [20418806](http://www.uniprot.org/citations/20418806)), PubMed: [23209295](http://www.uniprot.org/citations/23209295)),

PubMed:<a href="http://www.uniprot.org/citations/28017329" target="\_blank">28017329</a>). Plays a role in mitophagy which contributes to regulate mitochondrial quantity and quality by eliminating the mitochondria to a basal level to fulfill cellular energy requirements and preventing excess ROS production (PubMed:<a href="http://www.uniprot.org/citations/23209295" target="\_blank">23209295</a>, PubMed:<a href="http://www.uniprot.org/citations/28017329" target="\_blank">28017329</a>). In response to cellular stress and upon mitochondria fission, binds C-18 ceramides and anchors autophagolysosomes to outer mitochondrial membranes to eliminate damaged mitochondria (PubMed:<a href="http://www.uniprot.org/citations/22922758" target="\_blank">22922758</a>). While LC3s are involved in elongation of the phagophore membrane, the GABARAP/GATE-16 subfamily is essential for a later stage in autophagosome maturation (PubMed:<a href="http://www.uniprot.org/citations/20418806" target="\_blank">20418806</a>, PubMed:<a href="http://www.uniprot.org/citations/23209295" target="\_blank">23209295</a>, PubMed:<a href="http://www.uniprot.org/citations/28017329" target="\_blank">28017329</a>). Promotes primary ciliogenesis by removing OFD1 from centriolar satellites via the autophagic pathway (PubMed:<a href="http://www.uniprot.org/citations/24089205" target="\_blank">24089205</a>). Through its interaction with the reticulophagy receptor TEX264, participates in the remodeling of subdomains of the endoplasmic reticulum into autophagosomes upon nutrient stress, which then fuse with lysosomes for endoplasmic reticulum turnover (PubMed:<a href="http://www.uniprot.org/citations/31006537" target="\_blank">31006537</a>, PubMed:<a href="http://www.uniprot.org/citations/31006538" target="\_blank">31006538</a>). Upon nutrient stress, directly recruits cofactor JMY to the phagophore membrane surfaces and promotes JMY's actin nucleation activity and autophagosome biogenesis during autophagy (PubMed:<a href="http://www.uniprot.org/citations/30420355" target="\_blank">30420355</a>).

#### Cellular Location

Cytoplasmic vesicle, autophagosome membrane; Lipid-anchor Endomembrane system; Lipid-anchor Mitochondrion membrane; Lipid-anchor. Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:Q9CQV6}. Cytoplasmic vesicle. Note=LC3-II binds to the autophagic membranes. LC3-II localizes with the mitochondrial inner membrane during Parkin-mediated mitophagy (PubMed:28017329). Localizes also to discrete punctae along the ciliary axoneme

#### Tissue Location

Most abundant in heart, brain, skeletal muscle and testis. Little expression observed in liver


#### Anti-LC3B Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Anti-LC3B Antibody - Images






Western blot analysis of LC3B expression in Hela (A), NIH3T3 (B), rat brain (C) whole cell lysates.



Immunohistochemical analysis of LC3B staining in rat brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of LC3B staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

#### **Anti-LC3B Antibody - Background**

Recombinant protein corresponding to human LC3B.

**Anti-LC3B Antibody - Citations**

- [PDK1 inhibition reduces autophagy and cell senescence through the PI3K/AKT signalling pathway in a cigarette smoke mouse emphysema model](#)
- [ATG5 and ATG7 Expression Levels Are Reduced in Cutaneous Melanoma and Regulated by NRF1](#)