

MAGEA6 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6168a**Specification**

MAGEA6 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	P43360
Other Accession	NP_005354
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	34891
Antigen Region	214-243

MAGEA6 Antibody (C-term) - Additional Information**Gene ID** 4105**Other Names**

Melanoma-associated antigen 6, Cancer/testis antigen 16, CT16, MAGE-6 antigen, MAGE3B antigen, MAGEA6, MAGE6

Target/Specificity

This MAGEA6 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 214-243 amino acids from the C-terminal region of human MAGEA6.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MAGEA6 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

MAGEA6 Antibody (C-term) - Protein Information**Name** MAGEA6 {ECO:0000303|PubMed:31267705, ECO:0000312|HGNC:HGNC:6804}**Function** Activator of ubiquitin ligase activity of RING-type zinc finger-containing E3

ubiquitin-protein ligases that acts as a repressor of autophagy (PubMed:[17942928](#), PubMed:[20864041](#), PubMed:[31267705](#)). May enhance ubiquitin ligase activity of TRIM28 and stimulate p53/TP53 ubiquitination by TRIM28. Proposed to act through recruitment and/or stabilization of the Ubl-conjugating enzyme (E2) at the E3:substrate complex (PubMed:[17942928](#), PubMed:[20864041](#)). May play a role in tumor transformation or aspects of tumor progression (PubMed:[17942928](#), PubMed:[20864041](#)). In vitro promotes cell viability in melanoma cell lines (PubMed:[17942928](#)).

Tissue Location


Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes

MAGEA6 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MAGEA6 Antibody (C-term) - Images



The anti-MAGEA6 C-term Antibody (Cat.#AP6168a) is used in Western blot to detect MAGEA6 in A549 lysate.

MAGEA6 Antibody (C-term) - Background

MAGEA6 is a member of the MAGEA gene family. The members of this family have their entire coding sequences located in the last exon, and the encoded proteins show 50 to 80% sequence identity between each other. The promoters and first exons of the MAGEA genes show considerable variability, suggesting that the existence of this gene family enables the same function to be expressed under different transcriptional controls. The MAGEA genes are expressed at a high level in a number of tumors of various histologic types, and are silent in normal tissues with the exception of testis and placenta. The MAGEA genes are clustered on chromosome Xq28. They may

be implicated in some hereditary disorders, such as dyskeratosis congenita.

MAGEA6 Antibody (C-term) - References

Tatsumi, T., et al., Clin. Cancer Res. 9(3):947-954 (2003).
Tatsumi, T., et al., J. Exp. Med. 196(5):619-628 (2002).
Imai, Y., et al., Gene 160(2):287-290 (1995).
Rogner, U.C., et al., Genomics 29(3):725-731 (1995).
De Plaen, E., et al., Immunogenetics 40(5):360-369 (1994).