

p53 Antibody (T55)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6266d

Specification

p53 Antibody (T55) - Product Information

Application	IF, WB, IHC-P,E
Primary Accession	P04637
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	43653
Antigen Region	33-62

p53 Antibody (T55) - Additional Information

Gene ID 7157

Other Names

Cellular tumor antigen p53, Antigen NY-CO-13, Phosphoprotein p53, Tumor suppressor p53, TP53, P53

Target/Specificity

This p53 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 33-62 amino acids from human p53.

Dilution

IF~~1:10~50
WB~~1:2000
IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

p53 Antibody (T55) is for research use only and not for use in diagnostic or therapeutic procedures.

p53 Antibody (T55) - Protein Information

Name TP53

Synonyms P53

Function Multifunctional transcription factor that induces cell cycle arrest, DNA repair or apoptosis upon binding to its target DNA sequence (PubMed:[11025664](#), PubMed:[12524540](#), PubMed:[12810724](#), PubMed:[15186775](#), PubMed:[15340061](#), PubMed:[17317671](#), PubMed:[17349958](#), PubMed:[19556538](#), PubMed:[20673990](#), PubMed:[20959462](#), PubMed:[22726440](#), PubMed:[24051492](#), PubMed:[24652652](#), PubMed:[35618207](#), PubMed:[36634798](#), PubMed:[38653238](#), PubMed:[9840937](#)). Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis depending on the physiological circumstances and cell type (PubMed:[11025664](#), PubMed:[12524540](#), PubMed:[12810724](#), PubMed:[15186775](#), PubMed:[15340061](#), PubMed:[17189187](#), PubMed:[17317671](#), PubMed:[17349958](#), PubMed:[19556538](#), PubMed:[20673990](#), PubMed:[20959462](#), PubMed:[22726440](#), PubMed:[24051492](#), PubMed:[24652652](#), PubMed:[38653238](#), PubMed:[9840937](#)). Negatively regulates cell division by controlling expression of a set of genes required for this process (PubMed:[11025664](#), PubMed:[12524540](#), PubMed:[12810724](#), PubMed:[15186775](#), PubMed:[15340061](#), PubMed:[17317671](#), PubMed:[17349958](#), PubMed:[19556538](#), PubMed:[20673990](#), PubMed:[20959462](#), PubMed:[22726440](#), PubMed:[24051492](#), PubMed:[24652652](#), PubMed:[9840937](#)). One of the activated genes is an inhibitor of cyclin-dependent kinases. Apoptosis induction seems to be mediated either by stimulation of BAX and FAS antigen expression, or by repression of Bcl-2 expression (PubMed:[12524540](#), PubMed:[17189187](#)). Its pro-apoptotic activity is activated via its interaction with PPP1R13B/ASPP1 or TP53BP2/ASPP2 (PubMed:[12524540](#)). However, this activity is inhibited when the interaction with PPP1R13B/ASPP1 or TP53BP2/ASPP2 is displaced by PPP1R13L/iASPP (PubMed:[12524540](#)). In cooperation with mitochondrial PPIF is involved in activating oxidative stress-induced necrosis; the function is largely independent of transcription. Induces the transcription of long intergenic non-coding RNA p21 (lincRNA-p21) and lincRNA-Mkln1. LincRNA-p21 participates in TP53-dependent transcriptional repression leading to apoptosis and seems to have an effect on cell-cycle regulation. Implicated in Notch signaling cross-over. Prevents CDK7 kinase activity when associated to CAK complex in response to DNA damage, thus stopping cell cycle progression. Isoform 2 enhances the transactivation activity of isoform 1 from some but not all TP53-inducible promoters. Isoform 4 suppresses transactivation activity and impairs growth suppression mediated by isoform 1. Isoform 7 inhibits isoform 1-mediated apoptosis. Regulates the circadian clock by repressing CLOCK-BMAL1-mediated transcriptional activation of PER2 (PubMed:[24051492](#)).

Cellular Location

Cytoplasm. Nucleus. Nucleus, PML body. Endoplasmic reticulum. Mitochondrion matrix. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome Note=Recruited into PML bodies together with CHEK2 (PubMed:[12810724](#)) Translocates to mitochondria upon oxidative stress (PubMed:[22726440](#)) Translocates to mitochondria in response to mitomycin C treatment (PubMed:[27323408](#)). Competitive inhibition of TP53 interaction with HSPA9/MOT-2 by UBXN2A results in increased protein abundance and subsequent translocation of TP53 to the nucleus (PubMed:[24625977](#)) [Isoform 2]: Nucleus. Cytoplasm. Note=Localized mainly in the nucleus with minor staining in the cytoplasm [Isoform 4]: Nucleus. Cytoplasm. Note=Predominantly nuclear but translocates to the cytoplasm following cell stress [Isoform 8]: Nucleus. Cytoplasm. Note=Localized in both nucleus and cytoplasm in most cells. In some cells, forms foci in the nucleus that are different from nucleoli

Tissue Location


Ubiquitous. Isoforms are expressed in a wide range of normal tissues but in a tissue-dependent manner. Isoform 2 is expressed in most normal tissues but is not detected in brain, lung, prostate, muscle, fetal brain, spinal cord and fetal liver. Isoform 3 is expressed in most normal tissues but is not detected in lung, spleen, testis, fetal brain, spinal cord and fetal liver. Isoform 7 is expressed in most normal tissues but is not detected in prostate, uterus, skeletal muscle and breast. Isoform 8 is detected only in colon, bone marrow, testis, fetal brain and intestine. Isoform 9 is expressed in most normal tissues but is not detected in brain, heart, lung, fetal liver, salivary gland, breast or intestine

p53 Antibody (T55) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

p53 Antibody (T55) - Images




Confocal immunofluorescent analysis of p53 Antibody (T55)(Cat#AP6266d) with A2058 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). Actin filaments have been labeled with Alexa Fluor 555 phalloidin (red). DAPI was used to stain the cell nuclear (blue).



Western blot analysis of TP53 (arrow) using rabbit polyclonal p53 Antibody (T55) (Cat.#AP6266d).

293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the TP53 gene (Lane 2) (Origene Technologies).



All lanes : Anti-p53 Antibody (T55) at 1:2000 dilution Lane 1: 293 whole cell lysate Lane 2: A431 whole cell lysate Lane 3: HT-29 whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 44 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



p53 Antibody (T55) (Cat. #AP6266d) immunohistochemistry analysis in formalin fixed and paraffin embedded human skin tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of p53 Antibody (T55) for immunohistochemistry. Clinical relevance has not been evaluated.

p53 Antibody (T55) - Background

Tumor protein p53, a nuclear protein, plays an essential role in the regulation of cell cycle, specifically in the transition from G0 to G1. It is found in very low levels in normal cells, however, in a variety of transformed cell lines, it is expressed in high amounts, and believed to contribute to transformation and malignancy. p53 is a DNA-binding protein containing DNA-binding, oligomerization and transcription activation domains. It is postulated to bind as a tetramer to a p53-binding site and activate expression of downstream genes that inhibit growth and/or invasion, and thus function as a tumor suppressor. Mutants of p53 that frequently occur in a number of different human cancers fail to bind the consensus DNA binding site, and hence cause the loss of

tumor suppressor activity. Alterations of the TP53 gene occur not only as somatic mutations in human malignancies, but also as germline mutations in some cancer-prone families with Li-Fraumeni syndrome.

p53 Antibody (T55) - References

Blanchette, P., et al., Mol. Cell. Biol. 24(21):9619-9629 (2004).
Adachi, K., et al., Oncogene 23(47):7791-7798 (2004).
Zhang, Y., et al., J. Biol. Chem. 279(41):42545-42551 (2004).
Anazawa, Y., et al., Oncogene 23(46):7621-7627 (2004).
Montagnoli, A., et al., Cancer Res. 64(19):7110-7116 (2004).

p53 Antibody (T55) - Citations

- [An upward 9.4 T static magnetic field inhibits DNA synthesis and increases ROS-P53 to suppress lung cancer growth](#)