

HIRA Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6277c**Specification**

HIRA Antibody (Center) - Product Information

Application	WB,E
Primary Accession	P54198
Other Accession	Q8QFR2 , Q61666 , P79987
Reactivity	Human
Predicted	Chicken, Mouse, Xenopus
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	111835
Antigen Region	431-461

HIRA Antibody (Center) - Additional Information**Gene ID** 7290**Other Names**

Protein HIRA, TUP1-like enhancer of split protein 1, HIRA, DGCR1, HIR, TUPLE1

Target/Specificity

This HIRA antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 431-461 amino acids from the Central region of human HIRA.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

HIRA Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

HIRA Antibody (Center) - Protein Information**Name** HIRA**Synonyms** DGCR1, HIR, TUPLE1

Function Cooperates with ASF1A to promote replication-independent chromatin assembly. Required for the periodic repression of histone gene transcription during the cell cycle. Required for the formation of senescence-associated heterochromatin foci (SAHF) and efficient senescence-associated cell cycle exit.

Cellular Location

Nucleus, PML body. Note=Primarily, though not exclusively, localized to the nucleus. Localizes to PML bodies immediately prior to onset of senescence

Tissue Location


Expressed at high levels in kidney, pancreas and skeletal muscle and at lower levels in brain, heart, liver, lung, and placenta.

HIRA Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HIRA Antibody (Center) - Images



HIRA Antibody (Center) (Cat. #AP6277c) western blot analysis in HeLa cell line lysates (35ug/lane). This demonstrates the HIRA antibody detected the HIRA protein (arrow).

HIRA Antibody (Center) - Background

HIRA is a histone chaperone that preferentially places the variant histone H3.3 in nucleosomes. Orthologs of this protein in yeast, flies, and plants are necessary for the formation of transcriptionally silent heterochromatin. It plays an important role in the formation of the senescence-associated heterochromatin foci. These foci likely mediate the irreversible cell cycle

changes that occur in senescent cells. It is considered the primary candidate protein in some haploinsufficiency syndromes such as DiGeorge syndrome, and insufficient production of the gene may disrupt normal embryonic development.

HIRA Antibody (Center) - References

Zhang,R., Mol. Cell. Biol. 27 (6), 2343-2358 (2007)
Tang,Y., Nat. Struct. Mol. Biol. 13 (10), 921-929 (2006)