

Cleaved-MMP-10 (F99) Polyclonal Antibody
Catalog # AP63134**Specification**

Cleaved-MMP-10 (F99) Polyclonal Antibody - Product Information

Application	WB
Primary Accession	P09238
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

Cleaved-MMP-10 (F99) Polyclonal Antibody - Additional Information**Gene ID** 4319**Other Names**

MMP10; STMY2; Stromelysin-2; SL-2; Matrix metalloproteinase-10; MMP-10; Transin-2

Dilution

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Cleaved-MMP-10 (F99) Polyclonal Antibody - Protein Information**Name** MMP10**Synonyms** STMY2**Function**

Can degrade fibronectin, gelatins of type I, III, IV, and V; weakly collagens III, IV, and V. Activates procollagenase.

Cellular Location

Secreted, extracellular space, extracellular matrix



Cleaved-MMP-10 (F99) Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Cleaved-MMP-10 (F99) Polyclonal Antibody - Images





Cleaved-MMP-10 (F99) Polyclonal Antibody - Background

Can degrade fibronectin, gelatins of type I, III, IV, and V; weakly collagens III, IV, and V. Activates procollagenase.