

Cleaved-MMP-14 (Y112) Polyclonal Antibody
Catalog # AP63136**Specification****Cleaved-MMP-14 (Y112) Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	P50281
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

Cleaved-MMP-14 (Y112) Polyclonal Antibody - Additional Information

Gene ID 4323

Other Names

MMP14; Matrix metalloproteinase-14; MMP-14; MMP-X1; Membrane-type matrix metalloproteinase 1; MT-MMP 1; MTMMP1; Membrane-type-1 matrix metalloproteinase; MT1-MMP; MT1MMP

Dilution

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Cleaved-MMP-14 (Y112) Polyclonal Antibody - Protein Information

Name MMP14

Function

Endopeptidase that degrades various components of the extracellular matrix such as collagen (PubMed:[8015608](http://www.uniprot.org/citations/8015608)). Essential for pericellular collagenolysis and modeling of skeletal and extraskeletal connective tissues during development (By similarity). Activates progelatinase A/MMP2, thereby acting as a positive regulator of cell growth and migration (PubMed:[22065321](http://www.uniprot.org/citations/22065321), PubMed:[8015608](http://www.uniprot.org/citations/8015608)). Involved in the formation of the fibrovascular tissues in association with pro-MMP2 (PubMed:[12714657](http://www.uniprot.org/citations/12714657), PubMed:[22065321](http://www.uniprot.org/citations/22065321)). May be involved in actin cytoskeleton reorganization by cleaving PTK7 (PubMed:[20837484](http://www.uniprot.org/citations/20837484)). Acts as a regulator of Notch signaling by mediating cleavage and inhibition of DLL1 (PubMed:[21572390](http://www.uniprot.org/citations/21572390)). Cleaves ADGRB1 to release vasculostatin-40 which inhibits angiogenesis (PubMed:[21572390](http://www.uniprot.org/citations/21572390)).

<http://www.uniprot.org/citations/22330140> target="_blank">22330140). Acts as a negative regulator of the GDF15-GFRAL aversive response by mediating cleavage and inactivation of GFRAL (PubMed:35177851).

Cellular Location

Cell membrane; Single-pass type I membrane protein. Melanosome. Cytoplasm Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065). Forms a complex with BST2 and localizes to the cytoplasm (PubMed:17081065)

Tissue Location


Expressed in stromal cells of colon, breast, and head and neck. Expressed in lung tumors.

Cleaved-MMP-14 (Y112) Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Cleaved-MMP-14 (Y112) Polyclonal Antibody - Images



Cleaved-MMP-14 (Y112) Polyclonal Antibody - Background

Endopeptidase that degrades various components of the extracellular matrix such as collagen. Activates progelatinase A. Essential for pericellular collagenolysis and modeling of skeletal and extraskeletal connective tissues during development (By similarity). May be involved in actin cytoskeleton reorganization by cleaving PTK7 (PubMed:20837484). Acts as a positive regulator of cell growth and migration via activation of MMP15. Involved in the formation of the fibrovascular

tissues in association with pro-MMP2 (PubMed:12714657). Cleaves ADGRB1 to release vasculostatin-40 which inhibits angiogenesis (PubMed:22330140).