

**Cleaved-Thrombin APII (R327) Polyclonal Antibody**  
Catalog # AP63152**Specification**

---

**Cleaved-Thrombin APII (R327) Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P00734</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

**Cleaved-Thrombin APII (R327) Polyclonal Antibody - Additional Information****Gene ID** 2147**Other Names**

F2; Prothrombin; Coagulation factor II

**Dilution**

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**Cleaved-Thrombin APII (R327) Polyclonal Antibody - Protein Information****Name** F2**Function**

Thrombin, which cleaves bonds after Arg and Lys, converts fibrinogen to fibrin and activates factors V, VII, VIII, XIII, and, in complex with thrombomodulin, protein C. Functions in blood homeostasis, inflammation and wound healing.

**Cellular Location**

Secreted, extracellular space.

**Tissue Location**


Expressed by the liver and secreted in plasma.

**Cleaved-Thrombin APII (R327) Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **Cleaved-Thrombin APII (R327) Polyclonal Antibody - Images**



### **Cleaved-Thrombin APII (R327) Polyclonal Antibody - Background**

Thrombin, which cleaves bonds after Arg and Lys, converts fibrinogen to fibrin and activates factors V, VII, VIII, XIII, and, in complex with thrombomodulin, protein C. Functions in blood homeostasis, inflammation and wound healing.