

E2F-1 (Acetyl-Lys120) Polyclonal Antibody
Catalog # AP63271**Specification**

E2F-1 (Acetyl-Lys120) Polyclonal Antibody - Product Information

| | |
|-------------------|--------------------------|
| Application | WB |
| Primary Accession | Q01094 |
| Reactivity | Human, Mouse, Rat |
| Host | Rabbit |
| Clonality | Polyclonal |

E2F-1 (Acetyl-Lys120) Polyclonal Antibody - Additional Information**Gene ID** 1869**Other Names**
E2F1 RBBP3**Dilution**
WB~~WB: 1:500-10000 ELISA: 1:10000**Format**
PBS, pH 7.4, containing 0.09% (W/V) sodium azide as Preservative and 50% Glycerol.**Storage Conditions**
-20°C**E2F-1 (Acetyl-Lys120) Polyclonal Antibody - Protein Information****Name** E2F1 {ECO:0000303|PubMed:8964493, ECO:0000312|HGNC:HGNC:3113}**Function**

Transcription activator that binds DNA cooperatively with DP proteins through the E2 recognition site, 5'-TTTC[CG]CGC-3' found in the promoter region of a number of genes whose products are involved in cell cycle regulation or in DNA replication (PubMed:10675335, PubMed:12717439, PubMed:17050006, PubMed:17704056, PubMed:18625225, PubMed:28992046). The DRTF1/E2F complex functions in the control of cell-cycle progression from G1 to S phase (PubMed:10675335, PubMed:12717439, PubMed:17704056). E2F1 binds preferentially RB1 in a cell-cycle dependent manner (PubMed:10675335, PubMed:12717439, PubMed:<a

<http://www.uniprot.org/citations/17704056> target="_blank">17704056). It can mediate both cell proliferation and TP53/p53- dependent apoptosis (PubMed:8170954). Blocks adipocyte differentiation by binding to specific promoters repressing CEBPA binding to its target gene promoters (PubMed:20176812). Directly activates transcription of PEG10 (PubMed:17050006, PubMed:18625225, PubMed:28992046). Positively regulates transcription of RRP1B (PubMed:20040599).

Cellular Location


Nucleus

E2F-1 (Acetyl-Lys120) Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

E2F-1 (Acetyl-Lys120) Polyclonal Antibody - Images



E2F-1 (Acetyl-Lys120) Polyclonal Antibody - Background

Transcription activator that binds DNA cooperatively with DP proteins through the E2 recognition site, 5'-TTTC[CG]CGC- 3' found in the promoter region of a number of genes whose products are involved in cell cycle regulation or in DNA replication. The DRTF1/E2F complex functions in the

control of cell-cycle progression from G1 to S phase. E2F1 binds preferentially RB1 in a cell-cycle dependent manner. It can mediate both cell proliferation and TP53/p53-dependent apoptosis. Blocks adipocyte differentiation by binding to specific promoters repressing CEBPA binding to its target gene promoters (PubMed:20176812). Positively regulates transcription of RRP1B (PubMed:20040599).