

E2F-1 (Acetyl-K125) Polyclonal Antibody
Catalog # AP63285**Specification****E2F-1 (Acetyl-K125) Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	Q01094
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

E2F-1 (Acetyl-K125) Polyclonal Antibody - Additional Information

Gene ID 1869

Other Names

Transcription factor E2F1 (E2F-1) (PBR3) (Retinoblastoma-associated protein 1) (RBAP-1) (Retinoblastoma-binding protein 3) (RBBP-3) (pRB-binding protein E2F-1)

Dilution

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

E2F-1 (Acetyl-K125) Polyclonal Antibody - Protein Information**Name** E2F1 {ECO:0000303|PubMed:8964493, ECO:0000312|HGNC:HGNC:3113}**Function**

Transcription activator that binds DNA cooperatively with DP proteins through the E2 recognition site, 5'-TTTC[CG]CGC-3' found in the promoter region of a number of genes whose products are involved in cell cycle regulation or in DNA replication (PubMed:10675335, PubMed:12717439, PubMed:17050006, PubMed:17704056, PubMed:18625225, PubMed:28992046). The DRTF1/E2F complex functions in the control of cell-cycle progression from G1 to S phase (PubMed:10675335, PubMed:12717439, PubMed:17704056). E2F1 binds preferentially RB1 in a cell-cycle dependent manner (PubMed:10675335, PubMed:10675335, PubMed:10675335).

<http://www.uniprot.org/citations/12717439> target="_blank">12717439, PubMed:17704056). It can mediate both cell proliferation and TP53/p53- dependent apoptosis (PubMed:8170954). Blocks adipocyte differentiation by binding to specific promoters repressing CEBPA binding to its target gene promoters (PubMed:20176812). Directly activates transcription of PEG10 (PubMed:17050006, PubMed:18625225, PubMed:28992046). Positively regulates transcription of RRP1B (PubMed:20040599).

Cellular Location


Nucleus

E2F-1 (Acetyl-K125) Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

E2F-1 (Acetyl-K125) Polyclonal Antibody - Images



E2F-1 (Acetyl-K125) Polyclonal Antibody - Background

Transcription activator that binds DNA cooperatively with DP proteins through the E2 recognition site, 5'-TTTC[CG]CGC- 3' found in the promoter region of a number of genes whose products are

involved in cell cycle regulation or in DNA replication. The DRTF1/E2F complex functions in the control of cell-cycle progression from G1 to S phase. E2F1 binds preferentially RB1 in a cell-cycle dependent manner. It can mediate both cell proliferation and TP53/p53-dependent apoptosis. Blocks adipocyte differentiation by binding to specific promoters repressing CEBPA binding to its target gene promoters (PubMed:20176812). Positively regulates transcription of RRP1B (PubMed:20040599).