

Histone H2A.Z (Acetyl Lys7) Polyclonal Antibody
Catalog # AP63444**Specification**

Histone H2A.Z (Acetyl Lys7) Polyclonal Antibody - Product Information

Application	WB
Primary Accession	POC055
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

Histone H2A.Z (Acetyl Lys7) Polyclonal Antibody - Additional Information**Gene ID** 3015**Other Names**

H2AFZ; H2AZ; Histone H2A.Z; H2A/z

Dilution

WB~~WB: 1:500-1000

Format

PBS, pH 7.4, containing 0.09% (W/V) sodium azide as Preservative and 50% Glycerol.

Storage Conditions

-20°C

Histone H2A.Z (Acetyl Lys7) Polyclonal Antibody - Protein Information**Name** H2AZ1 ([HGNC:4741](#))**Function**

Variant histone H2A which replaces conventional H2A in a subset of nucleosomes. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post- translational modifications of histones, also called histone code, and nucleosome remodeling. May be involved in the formation of constitutive heterochromatin. May be required for chromosome segregation during cell division.

Cellular Location


Nucleus. Chromosome.

Histone H2A.Z (Acetyl Lys7) Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Histone H2A.Z (Acetyl Lys7) Polyclonal Antibody - Images



Histone H2A.Z (Acetyl Lys7) Polyclonal Antibody - Background

Variant histone H2A which replaces conventional H2A in a subset of nucleosomes. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. May be involved in the formation of constitutive heterochromatin. May be required for chromosome segregation during cell division.