

VE-Cadherin mouse Monoclonal Antibody(8E5)
Catalog # AP63782**Specification****VE-Cadherin mouse Monoclonal Antibody(8E5) - Product Information**

Application	IHC
Primary Accession	P33151
Reactivity	Human, Rat, Mouse
Host	Mouse
Clonality	Monoclonal

VE-Cadherin mouse Monoclonal Antibody(8E5) - Additional Information**Gene ID** 1003**Other Names**
CDH5**Dilution**
IHC~~IHC 1:100-200**Format**
Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.**Storage Conditions**
-20°C**VE-Cadherin mouse Monoclonal Antibody(8E5) - Protein Information****Name** CDH5 ([HGNC:1764](#))**Function**

Cadherins are calcium-dependent cell adhesion proteins (By similarity). They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types (PubMed:[21269602](http://www.uniprot.org/citations/21269602)). This cadherin may play an important role in endothelial cell biology through control of the cohesion and organization of the intercellular junctions (By similarity). It associates with alpha-catenin forming a link to the cytoskeleton (PubMed:[10861224](http://www.uniprot.org/citations/10861224)). Acts in concert with KRIT1 and PALS1 to establish and maintain correct endothelial cell polarity and vascular lumen (By similarity). These effects are mediated by recruitment and activation of the Par polarity complex and RAP1B (PubMed:[20332120](http://www.uniprot.org/citations/20332120)). Required for activation of PRKCZ and for the localization of phosphorylated PRKCZ, PARD3, TIAM1 and RAP1B to the cell junction (PubMed:[20332120](http://www.uniprot.org/citations/20332120)).

Cellular Location

Cell junction, adherens junction. Cell membrane; Single-pass type I membrane protein
Note=Found at cell-cell boundaries and probably at cell-matrix boundaries. KRIT1 and CDH5 reciprocally regulate their localization to endothelial cell-cell junctions.

Tissue Location



Endothelial tissues and brain.

VE-Cadherin mouse Monoclonal Antibody(8E5) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

VE-Cadherin mouse Monoclonal Antibody(8E5) - Images



VE-Cadherin mouse Monoclonal Antibody(8E5) - Background

Cadherins are calcium-dependent cell adhesion proteins. They preferentially interact with

themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types. This cadherin may play an important role in endothelial cell biology through control of the cohesion and organization of the intercellular junctions. It associates with alpha-catenin forming a link to the cytoskeleton. Acts in concert with KRIT1 to establish and maintain correct endothelial cell polarity and vascular lumen. These effects are mediated by recruitment and activation of the Par polarity complex and RAP1B. Required for activation of PRKCZ and for the localization of phosphorylated PRKCZ, PARD3, TIAM1 and RAP1B to the cell junction.