

CDC23 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6613a**Specification**

CDC23 Antibody (N-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	O9UJX2
Other Accession	A1A4R8
Reactivity	Human
Predicted	Bovine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	68834
Antigen Region	140-168

CDC23 Antibody (N-term) - Additional Information**Gene ID** 8697**Other Names**

Cell division cycle protein 23 homolog, Anaphase-promoting complex subunit 8, APC8, Cyclosome subunit 8, CDC23, ANAPC8

Target/Specificity

This CDC23 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 140-168 amino acids from the N-terminal region of human CDC23.

DilutionWB~~1:1000
IHC-P~~1:10~50**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CDC23 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CDC23 Antibody (N-term) - Protein Information**Name** CDC23

Synonyms ANAPC8


Function Component of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle (PubMed:[18485873](#)). The APC/C complex acts by mediating ubiquitination and subsequent degradation of target proteins: it mainly mediates the formation of 'Lys-11'-linked polyubiquitin chains and, to a lower extent, the formation of 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains (PubMed:[18485873](#)). The APC/C complex catalyzes assembly of branched 'Lys-11'-/'Lys-48'-linked branched ubiquitin chains on target proteins (PubMed:[29033132](#)).

CDC23 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CDC23 Antibody (N-term) - Images



Western blot analysis of CDC23 antibody (N-term) (Cat. #AP6613a) in HL60 cell line lysates(35ug/lane). CDC23 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human brain tissue reacted with CDC23 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

CDC23 Antibody (N-term) - Background

CDC23 shares strong similarity with *Saccharomyces cerevisiae* Cdc23, a protein essential for cell cycle progression through the G2/M transition. This protein is a component of anaphase-promoting complex (APC), which is composed of eight protein subunits and highly conserved in eukaryotic cells. APC catalyzes the formation of cyclin B-ubiquitin conjugate that is responsible for the ubiquitin-mediated proteolysis of B-type cyclins. This protein and 3 other members of the APC complex contain the TPR (tetratricopeptide repeat), a protein domain important for protein-protein interaction.

CDC23 Antibody (N-term) - References

Zhao,N., Genomics 53 (2), 184-190 (1998)