

PRELP Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6665C

Specification

PRELP Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	P51888
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	119-148

PRELP Antibody (Center) - Additional Information

Gene ID 5549

Other Names

Prolargin, Proline-arginine-rich end leucine-rich repeat protein, PRELP, SLRR2A

Target/Specificity

This PRELP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 119-148 amino acids from the Central region of human PRELP.

Dilution

WB~~1:1000
IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PRELP Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

PRELP Antibody (Center) - Protein Information

Name PRELP

Synonyms SLRR2A

Function May anchor basement membranes to the underlying connective tissue.

Cellular Location

Secreted, extracellular space, extracellular matrix


Tissue Location

Connective tissue.


PRELP Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PRELP Antibody (Center) - Images

PRELP Antibody (Center) (Cat. #AP6665c) western blot analysis in mouse heart tissue lysates (15ug/lane). This demonstrates the PRELP antibody detected PRELP protein (arrow).



PRELP Antibody (Center) (Cat. #AP6665c) IHC analysis in formalin fixed and paraffin embedded breast carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining.

This data demonstrates the use of the PRELP Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

PRELP Antibody (Center) - Background

PRELP is a leucine-rich repeat protein present in connective tissue extracellular matrix. This protein functions as a molecule anchoring basement membranes to the underlying connective tissue. This protein has been shown to bind type I collagen to basement membranes and type II collagen to cartilage. It also binds the basement membrane heparan sulfate proteoglycan perlecan. This protein is suggested to be involved in the pathogenesis of Hutchinson-Gilford progeria (HGP), which is reported to lack the binding of collagen in basement membranes and cartilage.

PRELP Antibody (Center) - References

Ehret, G.B., et al. Eur. J. Hum. Genet. 17(12):1650-1657(2009)
Grover, J., et al. Matrix Biol. 26(2):140-143(2007)
Johnson, J.M., et al. Mol. Vis. 12, 1057-1066 (2006)