

RAN Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6693b**Specification**

RAN Antibody (Center) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	P62826
Other Accession	P52301 , P62828 , P62827 , Q4R4M9 , Q9VZ23 , P79735 , P42558 , Q3T054 , NP_006316.1
Reactivity	Human, Mouse
Predicted	Bovine, Chicken, Zebrafish, Drosophila, Monkey, Rat, Xenopus
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	24423
Antigen Region	112-140

RAN Antibody (Center) - Additional Information**Gene ID** 5901**Other Names**

GTP-binding nuclear protein Ran, Androgen receptor-associated protein 24, GTPase Ran, Ras-like protein TC4, Ras-related nuclear protein, RAN, ARA24

Target/Specificity

This RAN antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 112-140 amino acids from the Central region of human RAN.

Dilution

WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RAN Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

RAN Antibody (Center) - Protein Information

Name RAN**Synonyms** ARA24 {ECO:0000303|PubMed:10400640}

Function GTPase involved in nucleocytoplasmic transport, participating both to the import and the export from the nucleus of proteins and RNAs (PubMed:[10400640](#), PubMed:[17209048](#), PubMed:[26272610](#), PubMed:[27306458](#), PubMed:[8276887](#), PubMed:[8636225](#), PubMed:[8692944](#), PubMed:[8896452](#), PubMed:[9351834](#), PubMed:[9428644](#), PubMed:[9822603](#)). Switches between a cytoplasmic GDP- and a nuclear GTP-bound state by nucleotide exchange and GTP hydrolysis (PubMed:[11336674](#), PubMed:[26272610](#), PubMed:[29040603](#), PubMed:[7819259](#), PubMed:[8636225](#), PubMed:[8692944](#), PubMed:[8896452](#), PubMed:[9351834](#), PubMed:[9428644](#), PubMed:[9822603](#)). Nuclear import receptors such as importin beta bind their substrates only in the absence of GTP-bound RAN and release them upon direct interaction with GTP-bound RAN, while export receptors behave in the opposite way. Thereby, RAN controls cargo loading and release by transport receptors in the proper compartment and ensures the directionality of the transport (PubMed:[8896452](#), PubMed:[9351834](#), PubMed:[9428644](#)). Interaction with RANBP1 induces a conformation change in the complex formed by XPO1 and RAN that triggers the release of the nuclear export signal of cargo proteins (PubMed:[20485264](#)). RAN (GTP-bound form) triggers microtubule assembly at mitotic chromosomes and is required for normal mitotic spindle assembly and chromosome segregation (PubMed:[10408446](#), PubMed:[29040603](#)). Required for normal progress through mitosis (PubMed:[12194828](#), PubMed:[29040603](#), PubMed:[8421051](#)). The complex with BIRC5/survivin plays a role in mitotic spindle formation by serving as a physical scaffold to help deliver the RAN effector molecule TPX2 to microtubules (PubMed:[18591255](#)). Acts as a negative regulator of the kinase activity of VRK1 and VRK2 (PubMed:[18617507](#)). Enhances AR-mediated transactivation. Transactivation decreases as the poly-Gln length within AR increases (PubMed:[10400640](#)).

Cellular Location

Nucleus. Nucleus envelope. Cytoplasm, cytosol Cytoplasm. Melanosome Note=Predominantly nuclear during interphase (PubMed:10679025, PubMed:12194828, PubMed:8421051). Becomes dispersed throughout the cytoplasm during mitosis (PubMed:12194828, PubMed:8421051). Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065).


Tissue Location


Expressed in a variety of tissues.

RAN Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


RAN Antibody (Center) - Images



RAN Antibody (Center) (Cat. #AP6693b) western blot analysis in mouse NIH-3T3 cell line lysates (35ug/lane). This demonstrates the RAN antibody detected the RAN protein (arrow).



RAN antibody (Center) (Cat. #AP6693b) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the RAN antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



RAN Antibody (Center) (Cat. #AP6693b) flow cytometric analysis of NIH-3T3 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

RAN Antibody (Center) - Background

RAN (ras-related nuclear protein) is a small GTP binding protein belonging to the RAS superfamily that is essential for the translocation of RNA and proteins through the nuclear pore complex. The RAN protein is also involved in control of DNA synthesis and cell cycle progression. Nuclear localization of RAN requires the presence of regulator of chromosome condensation 1 (RCC1). Mutations in RAN disrupt DNA synthesis. Because of its many functions, it is likely that RAN interacts with several other proteins. RAN regulates formation and organization of the microtubule network independently of its role in the nucleus-cytosol exchange of macromolecules. RAN could be a key signaling molecule regulating microtubule polymerization during mitosis. RCC1 generates a high local concentration of RAN-GTP around chromatin which, in turn, induces the local nucleation of microtubules. RAN is an androgen receptor (AR) coactivator that binds differentially with different lengths of polyglutamine within the androgen receptor. Polyglutamine repeat expansion in the AR is linked to Kennedy's disease (X-linked spinal and bulbar muscular atrophy). RAN coactivation of the AR diminishes with polyglutamine expansion within the AR, and this weak coactivation may lead to partial androgen insensitivity during the development of Kennedy's disease.

RAN Antibody (Center) - References

Kim, J.S., et al. Mol. Carcinog. 49(10):913-921(2010)
Liu, C., et al. Mol. Cell. Biol. 30(16):3956-3969(2010)
Boni, V., et al. Pharmacogenomics J. (2010) In press :
Ly, T.K., et al. J. Biol. Chem. 285(8):5815-5826(2010)
Mishra, R.K., et al. Nat. Cell Biol. 12(2):164-169(2010)