

**JAK1 (phospho Tyr1022) Polyclonal Antibody**  
Catalog # AP67086**Specification****JAK1 (phospho Tyr1022) Polyclonal Antibody - Product Information**

Application	IF
Primary Accession	<a href="#">P23458</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

**JAK1 (phospho Tyr1022) Polyclonal Antibody - Additional Information****Gene ID** 3716**Other Names**

JAK1; JAK1A; JAK1B; Tyrosine-protein kinase JAK1; Janus kinase 1; JAK-1

**Dilution**

IF~IF: 1:50-200 Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**JAK1 (phospho Tyr1022) Polyclonal Antibody - Protein Information****Name** JAK1**Synonyms** JAK1A, JAK1B**Function**

Tyrosine kinase of the non-receptor type, involved in the IFN-alpha/beta/gamma signal pathway (PubMed: [16239216](http://www.uniprot.org/citations/16239216)), PubMed: [28111307](http://www.uniprot.org/citations/28111307), PubMed: [32750333](http://www.uniprot.org/citations/32750333), PubMed: [7615558](http://www.uniprot.org/citations/7615558), PubMed: [8232552](http://www.uniprot.org/citations/8232552)). Kinase partner for the interleukin (IL)-2 receptor (PubMed: [11909529](http://www.uniprot.org/citations/11909529)) as well as interleukin (IL)-10 receptor (PubMed: [12133952](http://www.uniprot.org/citations/12133952)). Kinase partner for the type I interferon receptor IFNAR2 (PubMed: [16239216](http://www.uniprot.org/citations/16239216), PubMed: [28111307](http://www.uniprot.org/citations/28111307), PubMed: [32750333](http://www.uniprot.org/citations/32750333)),

PubMed:<a href="http://www.uniprot.org/citations/7615558" target="\_blank">7615558</a>, PubMed:<a href="http://www.uniprot.org/citations/8232552" target="\_blank">8232552</a>). In response to interferon-binding to IFNAR1-IFNAR2 heterodimer, phosphorylates and activates its binding partner IFNAR2, creating docking sites for STAT proteins (PubMed:<a href="http://www.uniprot.org/citations/7759950" target="\_blank">7759950</a>). Directly phosphorylates STAT proteins but also activates STAT signaling through the transactivation of other JAK kinases associated with signaling receptors (PubMed:<a href="http://www.uniprot.org/citations/16239216" target="\_blank">16239216</a>, PubMed:<a href="http://www.uniprot.org/citations/32750333" target="\_blank">32750333</a>, PubMed:<a href="http://www.uniprot.org/citations/8232552" target="\_blank">8232552</a>).

#### Cellular Location

Endomembrane system; Peripheral membrane protein. Note=Wholly intracellular, possibly membrane associated

#### Tissue Location



Expressed at higher levels in primary colon tumors than in normal colon tissue. The expression level in metastatic colon tumors is comparable to the expression level in normal colon tissue





### JAK1 (phospho Tyr1022) Polyclonal Antibody - Protocols





Provided below are standard protocols that you may find useful for product applications.





- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)





### JAK1 (phospho Tyr1022) Polyclonal Antibody - Images
















### JAK1 (phospho Tyr1022) Polyclonal Antibody - Background

Tyrosine kinase of the non-receptor type, involved in the IFN-alpha/beta/gamma signal pathway (PubMed:7615558). Kinase partner for the interleukin (IL)-2 receptor (PubMed:11909529) as well as interleukin (IL)-10 receptor (PubMed:12133952).