

Cyclin D1 (phospho Thr286) Polyclonal Antibody Catalog # AP67229

Specification

Cyclin D1 (phospho Thr286) Polyclonal Antibody - Product Information

Application	WB
Primary Accession	P24385
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

Cyclin D1 (phospho Thr286) Polyclonal Antibody - Additional Information

Gene ID 595

Other Names

CCND1; BCL1; PRAD1; G1/S-specific cyclin-D1; B-cell lymphoma 1 protein; BCL-1; BCL-1 oncogene; PRAD1 oncogene

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Cyclin D1 (phospho Thr286) Polyclonal Antibody - Protein Information

Name CCND1 {ECO:0000303|PubMed:8204893, ECO:0000312|HGNC:HGNC:1582}

Function

Regulatory component of the cyclin D1-CDK4 (DC) complex that phosphorylates and inhibits members of the retinoblastoma (RB) protein family including RB1 and regulates the cell-cycle during G(1)/S transition (PubMed:1827756, PubMed:1833066, PubMed:19412162, PubMed:33854235, PubMed:8114739, PubMed:8302605). Phosphorylation of RB1 allows dissociation of the transcription factor E2F from the RB/E2F complex and the subsequent transcription of E2F target genes which are responsible for the progression through the G(1) phase (PubMed:1827756, PubMed:1833066, PubMed:19412162, PubMed:33854235, PubMed:8114739, PubMed:8302605).

[8114739](http://www.uniprot.org/citations/8114739), PubMed:<[8302605](http://www.uniprot.org/citations/8302605)>). Hypophosphorylates RB1 in early G(1) phase (PubMed:<[1827756](http://www.uniprot.org/citations/1827756)>, PubMed:<[1833066](http://www.uniprot.org/citations/1833066)>, PubMed:<[19412162](http://www.uniprot.org/citations/19412162)>, PubMed:<[8114739](http://www.uniprot.org/citations/8114739)>, PubMed:<[8302605](http://www.uniprot.org/citations/8302605)>). Cyclin D-CDK4 complexes are major integrators of various mitogenic and antimitogenic signals (PubMed:<[1827756](http://www.uniprot.org/citations/1827756)>, PubMed:<[1833066](http://www.uniprot.org/citations/1833066)>, PubMed:<[19412162](http://www.uniprot.org/citations/19412162)>, PubMed:<[8302605](http://www.uniprot.org/citations/8302605)>). Also a substrate for SMAD3, phosphorylating SMAD3 in a cell-cycle-dependent manner and repressing its transcriptional activity (PubMed:<[15241418](http://www.uniprot.org/citations/15241418)>). Component of the ternary complex, cyclin D1/CDK4/CDKN1B, required for nuclear translocation and activity of the cyclin D-CDK4 complex (PubMed:<[9106657](http://www.uniprot.org/citations/9106657)>). Exhibits transcriptional corepressor activity with INSM1 on the NEUROD1 and INS promoters in a cell cycle-independent manner (PubMed:<[16569215](http://www.uniprot.org/citations/16569215)>, PubMed:<[18417529](http://www.uniprot.org/citations/18417529)>).

Cellular Location


Nucleus. Cytoplasm Nucleus membrane. Note=Cyclin D-CDK4 complexes accumulate at the nuclear membrane and are then translocated to the nucleus through interaction with KIP/CIP family members



Cyclin D1 (phospho Thr286) Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Cyclin D1 (phospho Thr286) Polyclonal Antibody - Images





Cyclin D1 (phospho Thr286) Polyclonal Antibody - Background

Regulatory component of the cyclin D1-CDK4 (DC) complex that phosphorylates and inhibits members of the retinoblastoma (RB) protein family including RB1 and regulates the cell-cycle during G(1)/S transition. Phosphorylation of RB1 allows dissociation of the transcription factor E2F from the RB/E2F complex and the subsequent transcription of E2F target genes which are responsible for the progression through the G(1) phase. Hypophosphorylates RB1 in early G(1) phase. Cyclin D-CDK4 complexes are major integrators of various mitogenic and antimitogenic signals. Also substrate for SMAD3, phosphorylating SMAD3 in a cell-cycle-dependent manner and repressing its transcriptional activity. Component of the ternary complex, cyclin D1/CDK4/CDKN1B, required for nuclear translocation and activity of the cyclin D-CDK4 complex. Exhibits transcriptional corepressor activity with INSM1 on the NEUROD1 and INS promoters in a cell cycle-independent manner.