

GRK 2 (phospho Ser685) Polyclonal Antibody
Catalog # AP67388**Specification****GRK 2 (phospho Ser685) Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	P25098
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

GRK 2 (phospho Ser685) Polyclonal Antibody - Additional Information**Gene ID** 156**Other Names**

ADRBK1; BARK; BARK1; GRK2; Beta-adrenergic receptor kinase 1; Beta-ARK-1; G-protein coupled receptor kinase 2

Dilution

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

GRK 2 (phospho Ser685) Polyclonal Antibody - Protein Information**Name** GRK2 ([HGNC:289](#))**Synonyms** ADRBK1, BARK, BARK1**Function**

Specifically phosphorylates the agonist-occupied form of the beta-adrenergic and closely related receptors, probably inducing a desensitization of them (PubMed: [19715378](http://www.uniprot.org/citations/19715378)). Key regulator of LPAR1 signaling (PubMed: [19306925](http://www.uniprot.org/citations/19306925)). Competes with RALA for binding to LPAR1 thus affecting the signaling properties of the receptor (PubMed: [19306925](http://www.uniprot.org/citations/19306925)). Desensitizes LPAR1 and LPAR2 in a phosphorylation-independent manner (PubMed: [19306925](http://www.uniprot.org/citations/19306925)). Positively regulates ciliary smoothened (SMO)-dependent Hedgehog (Hh) signaling pathway by facilitating the trafficking of SMO into the cilium and the stimulation of SMO activity (By similarity). Inhibits relaxation of airway smooth muscle in response to blue light (PubMed: [30284927](http://www.uniprot.org/citations/30284927)).

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:P26817}. Cell membrane {ECO:0000250|UniProtKB:P21146}. Postsynapse {ECO:0000250|UniProtKB:P26817}. Presynapse {ECO:0000250|UniProtKB:P26817}

Tissue Location


Expressed in peripheral blood leukocytes.

GRK 2 (phospho Ser685) Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GRK 2 (phospho Ser685) Polyclonal Antibody - Images



GRK 2 (phospho Ser685) Polyclonal Antibody - Background

Specifically phosphorylates the agonist-occupied form of the beta-adrenergic and closely related receptors, probably inducing a desensitization of them. Key regulator of LPAR1 signaling. Competes with RALA for binding to LPAR1 thus affecting the signaling properties of the receptor. Desensitizes LPAR1 and LPAR2 in a phosphorylation-independent manner (PubMed:19306925, PubMed:19715378). Positively regulates ciliary smoothened (SMO)- dependent Hedgehog (Hh) signaling pathway by facilitating the trafficking of SMO into the cilium and the stimulation of SMO activity (By similarity).