

**WAVE1 (phospho Tyr125) Polyclonal Antibody**  
Catalog # AP67612**Specification****WAVE1 (phospho Tyr125) Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q92558</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

**WAVE1 (phospho Tyr125) Polyclonal Antibody - Additional Information****Gene ID** 8936**Other Names**

WASF1; KIAA0269; SCAR1; WAVE1; Wiskott-Aldrich syndrome protein family member 1; WASP family protein member 1; Protein WAVE-1; Verprolin homology domain-containing protein 1

**Dilution**

WB~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**WAVE1 (phospho Tyr125) Polyclonal Antibody - Protein Information****Name** WASF1 ([HGNC:12732](#))**Synonyms** KIAA0269, SCAR1, WAVE1**Function**

Downstream effector molecule involved in the transmission of signals from tyrosine kinase receptors and small GTPases to the actin cytoskeleton. Promotes formation of actin filaments. Part of the WAVE complex that regulates lamellipodia formation (PubMed:[29961568](http://www.uniprot.org/citations/29961568)). The WAVE complex regulates actin filament reorganization via its interaction with the Arp2/3 complex (By similarity). As component of the WAVE1 complex, required for BDNF-NTRK2 endocytic trafficking and signaling from early endosomes (By similarity). Also involved in the regulation of mitochondrial dynamics (PubMed:[29961568](http://www.uniprot.org/citations/29961568)).

**Cellular Location**

Cytoplasm, cytoskeleton. Synapse {ECO:0000250|UniProtKB:Q5BJU7} Cell junction, focal

adhesion. Note=Dot- like pattern in the cytoplasm. Concentrated in Rac-regulated membrane-ruffling areas (PubMed:9889097). Partial translocation to focal adhesion sites might be mediated by interaction with SORBS2 (PubMed:18559503). In neurons, colocalizes with activated NTRK2 after BDNF addition in endocytic sites through the association with TMEM108 (By similarity). {ECO:0000250|UniProtKB:Q8R5H6, ECO:0000269|PubMed:18559503, ECO:0000269|PubMed:9889097}

#### Tissue Location


Highly expressed in brain. Lowly expressed in testis, ovary, colon, kidney, pancreas, thymus, small intestine and peripheral blood


#### WAVE1 (phospho Tyr125) Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### WAVE1 (phospho Tyr125) Polyclonal Antibody - Images





### WAVE1 (phospho Tyr125) Polyclonal Antibody - Background

Downstream effector molecule involved in the transmission of signals from tyrosine kinase receptors and small GTPases to the actin cytoskeleton. Promotes formation of actin filaments. Part of the WAVE complex that regulates lamellipodia formation. The WAVE complex regulates actin filament reorganization via its interaction with the Arp2/3 complex (By similarity). As component of the WAVE1 complex, required for BDNF- NTRK2 endocytic trafficking and signaling from early endosomes (By similarity).