

PTPN11 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6856B

Specification

PTPN11 Antibody (C-term) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	Q06124
Other Accession	P41499 , P35235 , Q90687
Reactivity	Human, Mouse
Predicted	Chicken, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	68011
Antigen Region	520-547

PTPN11 Antibody (C-term) - Additional Information

Gene ID 5781

Other Names

Tyrosine-protein phosphatase non-receptor type 11, Protein-tyrosine phosphatase 1D, PTP-1D, Protein-tyrosine phosphatase 2C, PTP-2C, SH-PTP2, SHP-2, Shp2, SH-PTP3, PTPN11, PTP2C, SHPTP2

Target/Specificity

This PTPN11 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 520-547 amino acids from the C-terminal region of human PTPN11.

Dilution

WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PTPN11 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

PTPN11 Antibody (C-term) - Protein Information

Name PTPN11

Synonyms PTP2C, SHPTP2

Function Acts downstream of various receptor and cytoplasmic protein tyrosine kinases to participate in the signal transduction from the cell surface to the nucleus (PubMed:[10655584](#), PubMed:[14739280](#), PubMed:[18559669](#), PubMed:[18829466](#), PubMed:[26742426](#), PubMed:[28074573](#)). Positively regulates MAPK signal transduction pathway (PubMed:[28074573](#)). Dephosphorylates GAB1, ARHGAP35 and EGFR (PubMed:[28074573](#)). Dephosphorylates ROCK2 at 'Tyr-722' resulting in stimulation of its RhoA binding activity (PubMed:[18559669](#)). Dephosphorylates CDC73 (PubMed:[26742426](#)). Dephosphorylates SOX9 on tyrosine residues, leading to inactivate SOX9 and promote ossification (By similarity). Dephosphorylates tyrosine-phosphorylated NEDD9/CAS-L (PubMed:[19275884](#)).

Cellular Location

Cytoplasm. Nucleus

Tissue Location


Widely expressed, with highest levels in heart, brain, and skeletal muscle.

PTPN11 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


PTPN11 Antibody (C-term) - Images



Western blot analysis of PTPN11 Antibody (C-term) (Cat. #AP6856b) in mouse bladder tissue lysates (35ug/lane). PTPN11 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human skeletal muscle reacted with PTPN11 Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



PTPN11 Antibody (C-term) (Cat. #AP6856b) flow cytometric analysis of Jurkat cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

PTPN11 Antibody (C-term) - Background

PTPN11 is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP contains two tandem Src homology-2 domains, which function as phospho-tyrosine binding domains and mediate the interaction of this PTP with its substrates.

PTPN11 Antibody (C-term) - References

Rikova, K., et al., Cell 131 (6), 1190-1203 (2007)