

**SPINK5 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP6869A**

**Specification**

---

**SPINK5 Antibody (N-term) - Product Information**

Application	WB, IHC-P, FC,E
Primary Accession	<a href="#">O9NQ38</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	120714
Antigen Region	188-217

**SPINK5 Antibody (N-term) - Additional Information**

**Gene ID** 11005

**Other Names**

Serine protease inhibitor Kazal-type 5, Lympho-epithelial Kazal-type-related inhibitor, LEKTI, Hemofiltrate peptide HF6478, Hemofiltrate peptide HF7665, SPINK5

**Target/Specificity**

This SPINK5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 188-217 amino acids from the N-terminal region of human SPINK5.

**Dilution**

WB~~1:1000  
IHC-P~~1:10~50  
FC~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

SPINK5 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**SPINK5 Antibody (N-term) - Protein Information**

**Name** SPINK5

**Function** Serine protease inhibitor, probably important for the anti- inflammatory and/or antimicrobial protection of mucous epithelia. Contribute to the integrity and protective barrier function of the skin by regulating the activity of defense-activating and desquamation- involved proteases. Inhibits KLK5, it's major target, in a pH-dependent manner. Inhibits KLK7, KLK14 CASP14, and trypsin.

**Cellular Location**

Secreted.

**Tissue Location**


Highly expressed in the thymus and stratum corneum. Also found in the oral mucosa, parathyroid gland, Bartholin's glands, tonsils, and vaginal epithelium. Very low levels are detected in lung, kidney, and prostate.

**SPINK5 Antibody (N-term) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


**SPINK5 Antibody (N-term) - Images**



Western blot analysis of SPINK5 Antibody (N-term) (Cat. #AP6869a) in A2058 cell line lysates (35ug/lane). SPINK5 (arrow) was detected using the purified Pab.



SPINK5 Antibody (N-term) (RB18954) IHC analysis in formalin fixed and paraffin embedded human tonsil tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the SPINK5 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



SPINK5 Antibody (N-term) (Cat. #AP6869a) flow cytometric analysis of A2058 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

### **SPINK5 Antibody (N-term) - Background**

SPINK5 is a multidomain serine protease inhibitor that contains 15 potential inhibitory domains. The inhibitor may play a role in skin and hair morphogenesis and anti-inflammatory and/or antimicrobial protection of mucous epithelia. Mutations may result in Netherton syndrome, a disorder characterized by ichthyosis, defective cornification, and atopy.

### **SPINK5 Antibody (N-term) - References**

Nin, M., et al., J. Dermatol. Sci. 54 (1), 17-24 (2009)