

CD130 Polyclonal Antibody
Catalog # AP68904**Specification****CD130 Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	P40189
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal

CD130 Polyclonal Antibody - Additional Information

Gene ID 3572

Other Names

IL6ST; Interleukin-6 receptor subunit beta; IL-6 receptor subunit beta; IL-6R subunit beta; IL-6R-beta; IL-6RB; CDw130; Interleukin-6 signal transducer; Membrane glycoprotein 130; gp130; Oncostatin-M receptor subunit alpha; CD antigen CD130

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

CD130 Polyclonal Antibody - Protein InformationName IL6ST ([HGNC:6021](#))**Function**

Signal-transducing molecule (PubMed: [2261637](http://www.uniprot.org/citations/2261637)). The receptor systems for IL6, LIF, OSM, CNTF, IL11, CTF1 and BSF3 can utilize IL6ST for initiating signal transmission. Binding of IL6 to IL6R induces IL6ST homodimerization and formation of a high-affinity receptor complex, which activates the intracellular JAK-MAPK and JAK-STAT3 signaling pathways (PubMed: [19915009](http://www.uniprot.org/citations/19915009), PubMed: [2261637](http://www.uniprot.org/citations/2261637), PubMed: [23294003](http://www.uniprot.org/citations/23294003)). That causes phosphorylation of IL6ST tyrosine residues which in turn activates STAT3 (PubMed: [19915009](http://www.uniprot.org/citations/19915009), PubMed: [23294003](http://www.uniprot.org/citations/23294003), PubMed: [25731159](http://www.uniprot.org/citations/25731159)). In parallel, the IL6 signaling pathway induces the expression of two cytokine receptor signaling inhibitors,

SOCS1 and SOCS3, which inhibit JAK and terminate the activity of the IL6 signaling pathway as a negative feedback loop (By similarity). Also activates the yes- associated protein 1 (YAP) and NOTCH pathways to control inflammation- induced epithelial regeneration, independently of STAT3 (By similarity). Acts as a receptor for the neuroprotective peptide humanin as part of a complex with IL27RA/WSX1 and CNTFR (PubMed:19386761). Mediates signals which regulate immune response, hematopoiesis, pain control and bone metabolism (By similarity). Has a role in embryonic development (By similarity). Essential for survival of motor and sensory neurons and for differentiation of astrocytes (By similarity). Required for expression of TRPA1 in nociceptive neurons (By similarity). Required for the maintenance of PTH1R expression in the osteoblast lineage and for the stimulation of PTH-induced osteoblast differentiation (By similarity). Required for normal trabecular bone mass and cortical bone composition (By similarity).

Cellular Location

[Isoform 1]: Cell membrane; Single-pass type I membrane protein

Tissue Location


Found in all the tissues and cell lines examined (PubMed:2261637). Expression not restricted to IL6 responsive cells (PubMed:2261637).

CD130 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD130 Polyclonal Antibody - Images



CD130 Polyclonal Antibody - Background

Signal-transducing molecule. The receptor systems for IL6, LIF, OSM, CNTF, IL11, CTF1 and BSF3 can utilize IL6ST for initiating signal transmission. Binding of IL6 to IL6R induces IL6ST homodimerization and formation of a high-affinity receptor complex, which activates Janus kinases (PubMed:2261637). That causes phosphorylation of IL6ST tyrosine residues which in turn activates STAT3 (PubMed:19915009, PubMed:23294003). Mediates signals which regulate immune response, hematopoiesis, pain control and bone metabolism (By similarity). Has a role in embryonic development (By similarity). Does not bind IL6 (PubMed:2261637). Essential for survival of motor and sensory neurons and for differentiation of astrocytes (By similarity). Required for expression of TRPA1 in nociceptive neurons (By similarity). Required for the maintenance of PTH1R expression in the osteoblast lineage and for the stimulation of PTH-induced osteoblast differentiation (By similarity). Required for normal trabecular bone mass and cortical bone composition (By similarity).