

CD137L Polyclonal Antibody
Catalog # AP68906**Specification**

CD137L Polyclonal Antibody - Product Information

Application	WB
Primary Accession	P41273
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

CD137L Polyclonal Antibody - Additional Information**Gene ID** 8744**Other Names**

TNFSF9; Tumor necrosis factor ligand superfamily member 9; 4-1BB ligand; 4-1BBL

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

CD137L Polyclonal Antibody - Protein Information**Name** TNFSF9**Function**

Cytokine that binds to TNFRSF9. Induces the proliferation of activated peripheral blood T-cells. May have a role in activation- induced cell death (AICD). May play a role in cognate interactions between T-cells and B-cells/macrophages.

Cellular Location

Membrane; Single-pass type II membrane protein.

Tissue Location



Expressed in brain, placenta, lung, skeletal muscle and kidney



CD137L Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD137L Polyclonal Antibody - Images





CD137L Polyclonal Antibody - Background

Cytokine that binds to TNFRSF9. Induces the proliferation of activated peripheral blood T-cells. May have a role in activation-induced cell death (AICD). May play a role in cognate interactions between T-cells and B-cells/macrophages.