

**CD284 Polyclonal Antibody**  
Catalog # AP68927**Specification****CD284 Polyclonal Antibody - Product Information**

Application	IF
Primary Accession	<a href="#">O00206</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal

**CD284 Polyclonal Antibody - Additional Information**

Gene ID 7099

**Other Names**

TLR4; Toll-like receptor 4; hToll; CD antigen CD284

**Dilution**

IF~IF: 1:50-200 Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**CD284 Polyclonal Antibody - Protein Information**

Name TLR4

**Function**

Transmembrane receptor that functions as a pattern recognition receptor recognizing pathogen- and damage-associated molecular patterns (PAMPs and DAMPs) to induce innate immune responses via downstream signaling pathways (PubMed: [10835634](http://www.uniprot.org/citations/10835634), PubMed: [15809303](http://www.uniprot.org/citations/15809303), PubMed: [16622205](http://www.uniprot.org/citations/16622205), PubMed: [17292937](http://www.uniprot.org/citations/17292937), PubMed: [17478729](http://www.uniprot.org/citations/17478729), PubMed: [20037584](http://www.uniprot.org/citations/20037584), PubMed: [20711192](http://www.uniprot.org/citations/20711192), PubMed: [23880187](http://www.uniprot.org/citations/23880187), PubMed: [27022195](http://www.uniprot.org/citations/27022195), PubMed: [29038465](http://www.uniprot.org/citations/29038465)). At the plasma membrane, cooperates with LY96 to mediate the innate immune response to bacterial lipopolysaccharide (LPS) (PubMed: [27022195](http://www.uniprot.org/citations/27022195))

target="\_blank">27022195</a>). Also involved in LPS-independent inflammatory responses triggered by free fatty acids, such as palmitate, and Ni(2+) (PubMed:<a href="http://www.uniprot.org/citations/20711192" target="\_blank">20711192</a>). Mechanistically, acts via MYD88, TIRAP and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response (PubMed:<a href="http://www.uniprot.org/citations/10835634" target="\_blank">10835634</a>, PubMed:<a href="http://www.uniprot.org/citations/21393102" target="\_blank">21393102</a>, PubMed:<a href="http://www.uniprot.org/citations/27022195" target="\_blank">27022195</a>, PubMed:<a href="http://www.uniprot.org/citations/36945827" target="\_blank">36945827</a>, PubMed:<a href="http://www.uniprot.org/citations/9237759" target="\_blank">9237759</a>). Alternatively, CD14-mediated TLR4 internalization via endocytosis is associated with the initiation of a MYD88-independent signaling via the TICAM1-TBK1-IRF3 axis leading to type I interferon production (PubMed:<a href="http://www.uniprot.org/citations/14517278" target="\_blank">14517278</a>). In addition to the secretion of proinflammatory cytokines, initiates the activation of NLRP3 inflammasome and formation of a positive feedback loop between autophagy and NF-kappa-B signaling cascade (PubMed:<a href="http://www.uniprot.org/citations/32894580" target="\_blank">32894580</a>). In complex with TLR6, promotes inflammation in monocytes/macrophages by associating with TLR6 and the receptor CD86 (PubMed:<a href="http://www.uniprot.org/citations/23880187" target="\_blank">23880187</a>). Upon ligand binding, such as oxLDL or amyloid-beta 42, the TLR4:TLR6 complex is internalized and triggers inflammatory response, leading to NF-kappa-B-dependent production of CXCL1, CXCL2 and CCL9 cytokines, via MYD88 signaling pathway, and CCL5 cytokine, via TICAM1 signaling pathway (PubMed:<a href="http://www.uniprot.org/citations/23880187" target="\_blank">23880187</a>). In myeloid dendritic cells, vesicular stomatitis virus glycoprotein G but not LPS promotes the activation of IRF7, leading to type I IFN production in a CD14-dependent manner (PubMed:<a href="http://www.uniprot.org/citations/15265881" target="\_blank">15265881</a>, PubMed:<a href="http://www.uniprot.org/citations/23880187" target="\_blank">23880187</a>). Required for the migration- promoting effects of ZG16B/PAUF on pancreatic cancer cells.

#### Cellular Location

Cell membrane; Single-pass type I membrane protein. Early endosome. Cell projection, ruffle {ECO:0000250|UniProtKB:Q9QUK6}. Note=Upon complex formation with CD36 and TLR6, internalized through dynamin-dependent endocytosis (PubMed:20037584). Colocalizes with RFTN1 at cell membrane and then together with RFTN1 moves to endosomes, upon lipopolysaccharide stimulation. Co-localizes with ZG16B/PAUF at the cell membrane of pancreatic cancer cells (PubMed:36232715)

#### Tissue Location

Highly expressed in placenta, spleen and peripheral blood leukocytes (PubMed:9237759, PubMed:9435236). Detected in monocytes, macrophages, dendritic cells and several types of T-cells (PubMed:27022195, PubMed:9237759). Expressed in pancreatic cancer cells but not in normal pancreatic cells (at protein level) (PubMed:36232715).






#### CD284 Polyclonal Antibody - Protocols




Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

• [Cell Culture](#)

**CD284 Polyclonal Antibody - Images**





### CD284 Polyclonal Antibody - Background

Cooperates with LY96 and CD14 to mediate the innate immune response to bacterial lipopolysaccharide (LPS) (PubMed:27022195). Acts via MYD88, TIRAP and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response (PubMed:9237759, PubMed:10835634, PubMed:27022195). Also involved in LPS-independent inflammatory responses triggered by free fatty acids, such as palmitate, and Ni(2+). Responses triggered by Ni(2+) require

non-conserved histidines and are, therefore, species-specific (PubMed:20711192). Both *M.tuberculosis* HSP70 (dnaK) and HSP65 (groEL-2) act via this protein to stimulate NF-kappa-B expression (PubMed:15809303). In complex with TLR6, promotes sterile inflammation in monocytes/macrophages in response to oxidized low-density lipoprotein (oxLDL) or amyloid-beta 42. In this context, the initial signal is provided by oxLDL- or amyloid- beta 42-binding to CD36. This event induces the formation of a heterodimer of TLR4 and TLR6, which is rapidly internalized and triggers inflammatory response, leading to the NF-kappa-B- dependent production of CXCL1, CXCL2 and CCL9 cytokines, via MYD88 signaling pathway, and CCL5 cytokine, via TICAM1 signaling pathway, as well as IL1B secretion. Binds electronegative LDL (LDL(-)) and mediates the cytokine release induced by LDL(-) (PubMed:23880187). Stimulation of monocytes in vitro with *M.tuberculosis* PstS1 induces p38 MAPK and ERK1/2 activation primarily via TLR2, but also partially via this receptor (PubMed:16622205).