

Claudin-1 Polyclonal Antibody
Catalog # AP69125**Specification****Claudin-1 Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	O95832
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

Claudin-1 Polyclonal Antibody - Additional Information**Gene ID** 9076**Other Names**

CLDN1; CLD1; SEMP1; Claudin-1; Senescence-associated epithelial membrane protein

Dilution

WB~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Claudin-1 Polyclonal Antibody - Protein Information**Name** CLDN1**Synonyms** CLD1, SEMP1**Function**

Claudins function as major constituents of the tight junction complexes that regulate the permeability of epithelia. While some claudin family members play essential roles in the formation of impermeable barriers, others mediate the permeability to ions and small molecules. Often, several claudin family members are coexpressed and interact with each other, and this determines the overall permeability. CLDN1 is required to prevent the paracellular diffusion of small molecules through tight junctions in the epidermis and is required for the normal barrier function of the skin. Required for normal water homeostasis and to prevent excessive water loss through the skin, probably via an indirect effect on the expression levels of other proteins, since CLDN1 itself seems to be dispensable for water barrier formation in keratinocyte tight junctions (PubMed: 23407391).

Cellular Location

Cell junction, tight junction. Cell membrane; Multi-pass membrane protein. Basolateral cell

membrane Note=Associates with CD81 and the CLDN1-CD81 complex localizes to the basolateral cell membrane.

Tissue Location


Strongly expressed in liver and kidney. Expressed in heart, brain, spleen, lung and testis.


Claudin-1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Claudin-1 Polyclonal Antibody - Images





Claudin-1 Polyclonal Antibody - Background

Claudins function as major constituents of the tight junction complexes that regulate the permeability of epithelia. While some claudin family members play essential roles in the formation of impermeable barriers, others mediate the permeability to ions and small molecules. Often, several claudin family members are coexpressed and interact with each other, and this determines the overall permeability. CLDN1 is required to prevent the paracellular diffusion of small molecules through tight junctions in the epidermis and is required for the normal barrier function of the skin. Required for normal water homeostasis and to prevent excessive water loss through the skin, probably via an indirect effect on the expression levels of other proteins, since CLDN1 itself seems to be dispensable for water barrier formation in keratinocyte tight junctions (PubMed:23407391).