

c-Myc Polyclonal Antibody
Catalog # AP69176**Specification****c-Myc Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	P01106
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

c-Myc Polyclonal Antibody - Additional Information**Gene ID** 4609**Other Names**

MYC; BHLHE39; Myc proto-oncogene protein; Class E basic helix-loop-helix protein 39; bHLHe39; Proto-oncogene c-Myc; Transcription factor p64

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

c-Myc Polyclonal Antibody - Protein Information**Name** MYC**Synonyms** BHLHE39**Function**

Transcription factor that binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3' (PubMed: [24940000](http://www.uniprot.org/citations/24940000), PubMed: [25956029](http://www.uniprot.org/citations/25956029)). Activates the transcription of growth-related genes (PubMed: [24940000](http://www.uniprot.org/citations/24940000), PubMed: [25956029](http://www.uniprot.org/citations/25956029)). Binds to the VEGFA promoter, promoting VEGFA production and subsequent sprouting angiogenesis (PubMed: [24940000](http://www.uniprot.org/citations/24940000), PubMed: [25956029](http://www.uniprot.org/citations/25956029)). Regulator of somatic reprogramming, controls self-renewal of embryonic stem cells (By similarity). Functions with TAF6L to activate target gene expression through RNA polymerase II pause release (By similarity). Positively regulates transcription of HNRNPA1, HNRNPA2 and PTBP1 which in turn

regulate splicing of pyruvate kinase PKM by binding repressively to sequences flanking PKM exon 9, inhibiting exon 9 inclusion and resulting in exon 10 inclusion and production of the PKM M2 isoform (PubMed:20010808).

Cellular Location


Nucleus, nucleoplasm. Nucleus, nucleolus. Nucleus. Cytoplasm

c-Myc Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

c-Myc Polyclonal Antibody - Images



c-Myc Polyclonal Antibody - Background

Transcription factor that binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3'. Activates the transcription of growth-related genes. Binds to the VEGFA promoter, promoting VEGFA production and subsequent sprouting angiogenesis (PubMed:24940000).