

CPE Polyclonal Antibody
Catalog # AP69265**Specification**

CPE Polyclonal Antibody - Product Information

Application	WB
Primary Accession	P16870
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

CPE Polyclonal Antibody - Additional Information**Gene ID** 1363**Other Names**CPE; Carboxypeptidase E; CPE; Carboxypeptidase H; CPH; Enkephalin convertase;
Prohormone-processing carboxypeptidase**Dilution**

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

CPE Polyclonal Antibody - Protein Information**Name** CPE**Function**



Sorting receptor that directs prohormones to the regulated secretory pathway. Acts also as a prohormone processing enzyme in neuro/endocrine cells, removing dibasic residues from the C-terminal end of peptide hormone precursors after initial endoprotease cleavage.



Cellular Location[Isoform 1]: Cytoplasmic vesicle, secretory vesicle {ECO:0000250|UniProtKB:Q00493}.
Cytoplasmic vesicle, secretory vesicle membrane {ECO:0000250|UniProtKB:P15087}; Peripheral membrane protein {ECO:0000250|UniProtKB:P15087}. Secreted {ECO:0000250|UniProtKB:P15087}. Note=Associated with the secretory granule membrane through direct binding to lipid rafts in intragranular conditions. {ECO:0000250|UniProtKB:Q00493}**CPE Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CPE Polyclonal Antibody - Images





CPE Polyclonal Antibody - Background

Removes residual C-terminal Arg or Lys remaining after initial endoprotease cleavage during prohormone processing. Processes proinsulin.