

**Cyclophilin F Polyclonal Antibody**  
Catalog # AP69368**Specification****Cyclophilin F Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P30405</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

**Cyclophilin F Polyclonal Antibody - Additional Information****Gene ID** 10105**Other Names**

PPIF; CYP3; Peptidyl-prolyl cis-trans isomerase F; mitochondrial; PPIase F; Cyclophilin F; Rotamase F

**Dilution**

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**Cyclophilin F Polyclonal Antibody - Protein Information****Name** PPIF**Synonyms** CYP3**Function**

PPIase that catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and may therefore assist protein folding (PubMed: [20676357](http://www.uniprot.org/citations/20676357)). Involved in regulation of the mitochondrial permeability transition pore (mPTP) (PubMed: [26387735](http://www.uniprot.org/citations/26387735)). It is proposed that its association with the mPTP is masking a binding site for inhibiting inorganic phosphate (Pi) and promotes the open probability of the mPTP leading to apoptosis or necrosis; the requirement of the PPIase activity for this function is debated (PubMed: [26387735](http://www.uniprot.org/citations/26387735)). In cooperation with mitochondrial p53/TP53 is involved in activating oxidative stress-induced necrosis (PubMed: [22726440](http://www.uniprot.org/citations/22726440)). Involved in modulation of mitochondrial membrane F(1)F(0)

ATP synthase activity and regulation of mitochondrial matrix adenine nucleotide levels (By similarity). Has anti-apoptotic activity independently of mPTP and in cooperation with BCL2 inhibits cytochrome c-dependent apoptosis (PubMed: <http://www.uniprot.org/citations/19228691> target="\_blank">19228691</a>).

#### Cellular Location


Mitochondrion matrix


#### Cyclophilin F Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Cyclophilin F Polyclonal Antibody - Images





### Cyclophilin F Polyclonal Antibody - Background

PPIase that catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and may therefore assist protein folding (PubMed:20676357). Involved in regulation of the mitochondrial permeability transition pore (mPTP). It is proposed that its association with the mPTP is masking a binding site for inhibiting inorganic phosphate (Pi) and promotes the open probability of the mPTP leading to apoptosis or necrosis; the requirement of the PPIase activity for this function is debated. In cooperation with mitochondrial TP53 is involved in activating oxidative stress-induced necrosis. Involved in modulation of mitochondrial membrane F(1)F(0) ATP synthase activity and regulation of mitochondrial matrix adenine nucleotide levels. Has anti-apoptotic activity independently of mPTP and in cooperation with BCL2 inhibits cytochrome c-dependent apoptosis.