

CYP2W1 Polyclonal Antibody
Catalog # AP69408**Specification****CYP2W1 Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	Q8TAV3
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

CYP2W1 Polyclonal Antibody - Additional Information

Gene ID 54905

Other Names

CYP2W1; Cytochrome P450 2W1; CYP11W1

Dilution

WB~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

CYP2W1 Polyclonal Antibody - Protein Information**Name** CYP2W1 {ECO:0000303|PubMed:26936974, ECO:0000312|HGNC:HGNC:20243}**Function**

A cytochrome P450 monooxygenase that may play a role in retinoid and phospholipid metabolism (PubMed:22591743, PubMed:26936974). Catalyzes the hydroxylation of saturated carbon hydrogen bonds. Hydroxylates all trans-retinoic acid (atRA) to 4- hydroxyretinoate and may regulate atRA clearance. Other retinoids such as all-trans retinol and all-trans retinal are potential endogenous substrates (PubMed:26936974). Catalyzes both epoxidation of double bonds and hydroxylation of carbon hydrogen bonds of the fatty acyl chain of 1-acylphospholipids/2-lysophospholipids. Can metabolize various lysophospholipids classes including lysophosphatidylcholines (LPCs), lysophosphatidylinositols (LPIs), lysophosphatidylserines (LPSs), lysophosphatidylglycerols (LPGs), lysophosphatidylethanolamines (LPEs) and lysophosphatidic acids (LPAs) (PubMed:22591743). Has low or no activity toward 2-acylphospholipids/1-lysophospholipids, diacylphospholipids and free fatty acids (PubMed:22591743).

target="_blank">22591743, PubMed:26936974). May play a role in tumorigenesis by activating procarcinogens such as aflatoxin B1, polycyclic aromatic hydrocarbon dihydrodiols and aromatic amines (PubMed:16551781, PubMed:20805301, PubMed:24278521). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (CPR; NADPH-ferrihemoprotein reductase) (PubMed:22591743, PubMed:26936974).

Cellular Location

Endoplasmic reticulum lumen. Cell membrane. Microsome membrane. Note=About 8% are expressed on the cell surface.

Tissue Location


Very low levels are detected in fetal and adult tissues. Highly expressed in several tumor samples, in particular colon and adrenal tumors.

CYP2W1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CYP2W1 Polyclonal Antibody - Images



CYP2W1 Polyclonal Antibody - Background

Seems to have broad catalytic activity towards several chemicals, including polycyclic aromatic hydrocarbon dihydrodiols and aromatic amines (PubMed:16551781, PubMed:24278521). Active also in the metabolism of indoline substrates and is able to activate aflatoxin B1 into cytotoxic products (PubMed:20805301). Furthermore, it seems to be involved in the oxydation of lysophospholipids and fatty acids (PubMed:22591743).