

D4DR Polyclonal Antibody
Catalog # AP69461**Specification**

D4DR Polyclonal Antibody - Product Information

Application	WB
Primary Accession	P21917
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

D4DR Polyclonal Antibody - Additional Information**Gene ID** 1815**Other Names**

DRD4; D(4) dopamine receptor; D(2C) dopamine receptor; Dopamine D4 receptor

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

D4DR Polyclonal Antibody - Protein Information**Name** DRD4**Function**

Dopamine receptor responsible for neuronal signaling in the mesolimbic system of the brain, an area of the brain that regulates emotion and complex behavior. Activated by dopamine, but also by epinephrine and norepinephrine, and by numerous synthetic agonists and drugs (PubMed: [16423344](http://www.uniprot.org/citations/16423344), PubMed: [27659709](http://www.uniprot.org/citations/27659709), PubMed: [29051383](http://www.uniprot.org/citations/29051383), PubMed: [9003072](http://www.uniprot.org/citations/9003072)). Agonist binding triggers signaling via G proteins that inhibit adenylyl cyclase (PubMed: [16423344](http://www.uniprot.org/citations/16423344), PubMed: [27659709](http://www.uniprot.org/citations/27659709), PubMed: [29051383](http://www.uniprot.org/citations/29051383), PubMed: [7512953](http://www.uniprot.org/citations/7512953), PubMed: [7643093](http://www.uniprot.org/citations/7643093)). Modulates the circadian rhythm of contrast sensitivity by regulating the rhythmic expression of NPAS2 in the retinal ganglion cells (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein

Tissue Location


Highly expressed in retina. Detected at much lower levels in brain, in amygdala, thalamus, hypothalamus, cerebellum and pituitary.


D4DR Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

D4DR Polyclonal Antibody - Images





D4DR Polyclonal Antibody - Background

Dopamine receptor responsible for neuronal signaling in the mesolimbic system of the brain, an area of the brain that regulates emotion and complex behavior. Activated by dopamine, but also by epinephrine and norepinephrine, and by numerous synthetic agonists and drugs (PubMed:9003072, PubMed:16423344, PubMed:27659709, PubMed:29051383). Agonist binding triggers signaling via G proteins that inhibit adenylyl cyclase (PubMed:7512953, PubMed:7643093, PubMed:16423344, PubMed:27659709, PubMed:29051383). Modulates the circadian rhythm of contrast sensitivity by regulating the rhythmic expression of NPAS2 in the retinal ganglion cells (By similarity).