

**DNAM-1 Polyclonal Antibody**  
**Catalog # AP69564****Specification****DNAM-1 Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q15762</a>
Reactivity	Human, Mouse, Monkey
Host	Rabbit
Clonality	Polyclonal

**DNAM-1 Polyclonal Antibody - Additional Information****Gene ID** 10666**Other Names**

CD226; DNAM1; CD226 antigen; DNAX accessory molecule 1; DNAM-1; CD antigen CD226

**Dilution**

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**DNAM-1 Polyclonal Antibody - Protein Information****Name** CD226**Synonyms** DNAM1**Function**

Cell surface receptor that plays an important role in the immune system, particularly in intercellular adhesion, lymphocyte signaling, cytotoxicity and lymphokine secretion mediated by cytotoxic T-cells and NK cells (PubMed: <a href="http://www.uniprot.org/citations/8673704" target="\_blank">8673704</a>, PubMed: <a href="http://www.uniprot.org/citations/9712030" target="\_blank">9712030</a>). Functions as a costimulatory receptor upon recognition of target cells, such as virus- infected or tumor cells. Upon binding to its ligands PVR/CD155 or NECTIN2/CD112 on target cells, promotes the cytotoxic activity of NK cells and CTLs, enhancing their ability to kill these cells (PubMed: <a href="http://www.uniprot.org/citations/26755705" target="\_blank">26755705</a>, PubMed: <a href="http://www.uniprot.org/citations/31253644" target="\_blank">31253644</a>, PubMed: <a href="http://www.uniprot.org/citations/30591568" target="\_blank">30591568</a>). Mechanistically, phosphorylation by Src kinases such as LYN of FYN, enables binding to adapter GRB2, leading to activation of VAV1, PI3K and PLCG1. Promotes also activation of kinases ERK and AKT, as well as calcium fluxes (By similarity).

**Cellular Location**

Cell membrane; Single-pass type I membrane protein. Note=Localizes to lipid rafts.


**Tissue Location**




Expressed by peripheral blood T-lymphocytes.



**DNAM-1 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**DNAM-1 Polyclonal Antibody - Images**





### DNAM-1 Polyclonal Antibody - Background

Involved in intercellular adhesion, lymphocyte signaling, cytotoxicity and lymphokine secretion mediated by cytotoxic T-lymphocyte (CTL) and NK cell (PubMed:8673704). Cell surface receptor for NECTIN2. Upon ligand binding, stimulates T- cell proliferation and cytokine production, including that of IL2, IL5, IL10, IL13, and IFNG. Competes with PVRIG for NECTIN2-binding (PubMed:26755705).